

CA1  
MS61  
-54 N42

GOVT. PUBNS  
21

CANADA  
DEPARTMENT OF MINES AND TECHNICAL SURVEYS


MINES BRANCH  
INDUSTRIAL MINERALS DIVISION

INDUSTRIAL WATER RESOURCES OF CANADA

WATER SURVEY REPORT NO. 6

FRASER RIVER DRAINAGE BASIN, 1950-51

By  
J. F. J. Thomas



EDMOND CLOUTIER, C.M.G., O.A., D.S.P.  
QUEEN'S PRINTER AND CONTROLLER OF STATIONERY  
OTTAWA, 1984

Price 75 cents

No. 842

3 1761 11767559 5







CA1  
MS 61  
-54N42

CANADA  
DEPARTMENT OF MINES AND TECHNICAL SURVEYS

---

MINES BRANCH  
INDUSTRIAL MINERALS DIVISION

---

INDUSTRIAL WATER RESOURCES OF CANADA

WATER SURVEY REPORT NO. 6

FRASER RIVER DRAINAGE BASIN, 1950-51


By  
J. F. J. Thomas




---

Price, 75 cents

No. 842







## CONTENTS

	PAGE
Introduction.....	5
Fraser River drainage basin.....	5
Survey procedure.....	6
Analytical procedure.....	6
Part I—Surface waters of Fraser River drainage basin	
Discussion.....	8
Summary.....	9
Part II—Municipal water supplies within Fraser River drainage basin.	
Description of municipal waterworks systems within Fraser River drainage basin.....	58
Discussion.....	87
Summary .....	87
Appendix A—Sampling locations of surface waters...	89
Appendix B—Municipalities with organized water systems within Fraser River drainage basin.....	91

## TABLES

I—Tidal influence on Fraser River.....	7
II—Chemical quality of surface waters in Fraser River drainage basin.....	14
III—Chemical quality of municipal water supplies within Fraser River drainage basin.....	74
IV—Municipal water supplies in Fraser River drainage basin—summary of data on area, total population, and population served.....	86
V—Municipal water supplies in Fraser River drainage basin—summary of data on systems including source, treatment and hardness of waters..	86

## FIGURES

Fig. 1. Map showing drainage basins under study in Western Canada.....	4
Fig. 2. Map showing the location of surface water sampling stations and municipal water supplies (in pocket)	
Fig. 3. Chart showing variation in water hardness in Fraser River.....	10
Fig. 4. Relationship between mineral content and river flow, Fraser River at Mission City, B.C.....	11
Fig. 5. Relationship between mineral content and river flow, Thompson River at Kamloops, B.C.....	12
Fig. 6. Relationship between mineral content and river flow, Quesnel River near Quesnel, B.C.....	13






FIGURE 1. MAP SHOWING DRAINAGE BASINS UNDER STUDY IN WESTERN CANADA



# INDUSTRIAL WATER RESOURCES OF CANADA

## Chemical Quality of Surface and Municipal Water Supplies in the Fraser River Drainage Basin, 1950-51

### INTRODUCTION

This report is the sixth in a series on the chemical quality of surface and municipal water supplies available for industrial and domestic use in Canada. Report No. 1<sup>1</sup> outlines the scope and procedures used in the country-wide survey and discusses the interpretation of analytical results to be recorded in subsequent reports. Reports No. 2<sup>2</sup> and 3<sup>3</sup> cover the results of studies on the Ottawa River and Upper St. Lawrence River-Central Great Lakes drainage basins respectively.

This report and reports Nos. 4 and 5 cover the areas and drainage basins in British Columbia outlined in Figure I. It will be noted that these three reports cover all of British Columbia except the northern portions drained by the Yukon and Mackenzie River systems, which areas will be covered in Report No. 8, now in preparation.

The northern portion of the province, in particular the northern portion of Fraser River and western portion of Skeena River basins, is now the centre of considerable industrial activity and it is expected that expansion of both industry and agriculture in these areas, now relatively sparsely settled, will necessitate more detailed studies of water quality in the near future.

The method of presentation of data on this watershed is similar to that of previous reports and no attempt has been made to discuss in detail all the information recorded in this report or obtained during the survey. Part I tabulates the analytical results of daily, monthly and spot sampling of surface waters in the basin. Part II reports similar analytical data obtained on municipal waters within the basin and also includes information on the operation of most of the organized water systems.

The co-operation and assistance given to the writer by W. C. Warren, District Engineer, and other engineering personnel of the Water Resources Division, Dept. of Resources and Development, Vancouver, B.C. in selecting sampling locations and in supplying the data on river and lake stage and river discharge, used in this report, is gratefully acknowledged.

The co-operation of municipal officials and water works engineers who supplied the writer with information on their waterworks systems by correspondence or during visits to their communities, is also greatly appreciated.

### FRASER RIVER DRAINAGE BASIN

The Fraser River, 850 miles in length, with its many tributaries drains about 91,660 square miles, all of which is in British Columbia except for some 220 square miles in the United States, south of Sumas, B.C. This river and its larger tributaries provide the passes and valleys that permit crossing the mountainous terrain of British Columbia. The railways and Trans-Canada Highway follow the South Thompson (206 miles in length), the Thompson and the lower Fraser River to the sea. Travel from the earliest days into the central areas of British Columbia, in particular the Cariboo and north western portion of the province, has been along the Fraser River and its central plateau. The Canadian National Railway follows the upper Fraser River, its large tributary, the Nechako River (287 miles in length) and the Skeena River to the coast at Prince Rupert.

The Fraser River, rising in the Rocky Mountains just west of the source of the large North Saskatchewan and Athabasca Rivers, flows through almost all types of terrain found in British Columbia. It flows west skirting the Cariboo Range then south along the central plateau lying between the coastal range and Cariboo and Monashee mountain ranges, then south and west through the coastal range to the sea.

One of the most fertile and heavily-populated areas in British Columbia and indeed in all Canada has developed on the delta of the Fraser River. Here the rich alluvial soil, and a climate tempered by the Japanese current, with abundant rainfall, has given rise to intensive farming particularly of small fruits and vegetables. The abundant rainfall along the coastal range and plain has resulted in heavy forestation and led to the formation of a large lumbering industry with headquarters in the delta area.

<sup>1</sup> Industrial Water Resources of Canada (Water Survey Report No. 1): Scope, Procedure, and Interpretation of Survey Studies, Mines Branch Report No. 833, Dept. of Mines and Technical Surveys, Ottawa, 1952.

<sup>2</sup> Industrial Water Resources of Canada (Water Survey Report No. 2): Ottawa River Drainage Basin, Mines Branch Report No. 834, Dept. of Mines and Technical Surveys, Ottawa, 1952.

<sup>3</sup> Industrial Water Resources of Canada (Water Survey Report No. 3): Upper St. Lawrence River-Central Great Lakes Drainage Basin in Canada, Mines Branch Report No. 837, Dept. of Mines and Technical Surveys, Ottawa. (In press.)



Although rainfall is less abundant in the interior plateau of the Fraser Basin, particularly the Cariboo region, the country is ideal grazing land. The wide valleys and plateaus in the upper Fraser, Nechako, Stuart and South Thompson River basins are now being opened to farming and industry. As in other parts of the province the river valleys are extremely fertile but there are only two large areas in the entire province with great agricultural possibilities, the Peace River block lying in the drainage basin of the Mackenzie River, and the Stuart Lake district. Both these areas are rapidly being opened to farming and industry.

As with any large river system in a mountainous region, the rivers are the lifeblood of the country. It is such tributary rivers as the North Thompson, 210 miles in length, the Chilcotin, 146 miles in length, and the West Road River, 140 miles in length, that are permitting the settlement and opening up of this province, rich in natural resources and industrial potential.

In all its length Fraser River is a turbulent, silt-laden stream, for the most part too rapid for transportation. This is also true of most of the tributary rivers except those in the upper lake regions of South Thompson, Nechako and Stuart Rivers, or those flowing for the most part in plateau areas.

## SURVEY PROCEDURE

The methods of sampling and survey procedure employed in this investigation were in general similar to those outlined in previous reports and given in detail in Water Survey Report No. 1<sup>1</sup>. The Fraser River system was studied during 1950-51, at the same time studies were carried out on water quality in the Skeena River basin<sup>2</sup>. Twenty sampling stations were operated in this watershed, two daily stations, and eighteen monthly stations.

At the daily stations samples were collected each day into 16 ounce, pressure-sealed bottles which were shipped thrice monthly by the collector to the British Columbia Research Council at Vancouver. Here data regarding the daily water temperature, water level, etc., were recorded, each daily sample tested for specific conductance and a 10-day composite sample prepared. These composite samples were tested for pH, colour, turbidity, specific conductance and alkalinity and then shipped to the Mines Branch laboratory at Ottawa where a complete analysis was carried out.

The monthly samples were shipped directly by the collector to the laboratory in Ottawa. Whenever possible, samples were also obtained from these stations when the river was at high and low flow.

During the summer of 1950, most of the accessible portion of the basin was travelled with a mobile laboratory, and municipal waters and additional samples of river and lake waters were collected and field-tested.

## ANALYTICAL PROCEDURE

The methods of analyses and the method of reporting analytical results used in this survey are essentially those outlined in detail in Water Survey Report No. 1.

Until June 15, 1950, all samples received in the Ottawa laboratory were stored unopened in the dark until analyses could be started. It will be noted from Table II that storage time on these earlier samples was usually quite brief. After June 15, 1950, all samples received in the laboratory were immediately tested for pH, colour, turbidity, alkalinity, specific conductance, total hardness and, sometimes, chloride and calcium ion content. Previous experience had shown that these determinations are those normally affected by storage. However, storage time on these samples is still reported as the total time elapsing between sampling and the beginning of final analysis, even though waters after June 15th were usually tested for the unstable constituents within a much shorter period.

The tests carried out by the British Columbia Research Council on composite samples were all repeated in the Ottawa laboratory. A comparison of maximum and minimum individual differences and the arithmetical mean or average of all test results in each laboratory shows that, as in previous work in the two laboratories, the major differences are in the determinations for colour and turbidity. The individual differences in these determinations, particularly colour, was in some cases quite large, and was to be expected because of the turbid nature of many of the waters. Colour determined in the British Columbia Research Council laboratories is in many cases "apparent colour" due to turbidity of the waters whereas in the Ottawa laboratory colour was determined normally on the supernatant or settled sample. It is well known that storage of waters may cause bleaching or loss of colour and coagulation or settling of turbidity. However, even though storage time when "immediate testing" was carried out at Ottawa was on the average almost twice as long as when tests were made in British Columbia, the described survey procedure does give quite satisfactory agreement for pH and alkalinity, two important values that often show considerable changes on storage. The maximum variation in pH between the laboratories on any one sample was 0.5.

<sup>1</sup> Industrial Water Resources of Canada (Water Survey Report No. 1): Scope, Procedure, and Interpretation of Survey Studies, Mines Branch Report No. 833, Dept. of Mines and Technical Surveys, Ottawa, 1952.

<sup>2</sup> Industrial Water Resources of Canada (Water Survey Report No. 5): Skeena River Drainage Basin, Vancouver Island, and Coastal Areas of British Columbia, 1949-51, Mines Branch Report No. 839, Dept. of Mines and Technical Surveys, Ottawa, 1953.



## PART I

### SURFACE WATERS OF THE FRASER RIVER DRAINAGE BASIN

Daily samples were collected of Fraser River from the railway bridge at Mission City, and of Thompson River from the highway bridge at Kamloops, during the period February, 1950 to February, 1951. When field work was being carried out in 1950 it was found that sampling at the latter station was from the southern side or Kamloops side of the river and that in many cases the water was probably South Thompson River water. This location is just below the junction of North and South Thompson Rivers and indications were that, at least during much of the year, complete mixing of the river waters does not occur.

During the summer of 1950 additional samples of surface waters and municipal water supplies within the basin were collected. The locations of all surface water sampling points within this watershed are outlined in Appendix A and are shown on the map of the basin, Figure 2 (in map pocket). As in other basins in this province, various areas, in particular the northern and northeastern portion of the basin, were inaccessible by road and consequently several large tributary rivers were not studied. Since these rivers either have their source in the same general area or traverse the same type of terrain as nearby streams which were studied it is believed that the quality of their waters can be assumed similar in character. These inaccessible areas are only sparsely settled and domestic and industrial use of the river waters is at present practically nil.

Most samples collected during field work in the summer were tested immediately in the mobile laboratory for the constituents and properties that may change on storage. These field results are reported in Tables II and III in brackets beside the results found later in the laboratory. Repetition of these tests indicates changes in water due to storage and enables estimation of the quality of the water *in situ*.

Table I tabulates available information on the influence of tides on Fraser River water. No special study was made in this regard at this time and the data shown in Table I were supplied by the Water Resources Division, Department of Resources and Development.

TABLE I  
Tidal Influence—Fraser River

#### A. RISING TIDE

Sampling Location	River Discharge at Hope, B.C. (second-feet)	River Depth at Sampling (feet)	Sampling		High Tide		Low Tide		Chloride Content, p.p.m.		
			Date	Average Time	Time	Height† feet	Time	Height† feet	1/5*	3/5*	4/5*
Steveston.....	27,100	13.5	13/12/48	12.25	14.28	12.4	9.15	9.3	15.02	.....	85.86
	128,000	8.0	26/ 4/49	13.21	17.33	12.2	11.15	5.0	4.9	.....	4.5
	318,000	9.0	17/ 5/49	19.10	23.19	14.7	15.22	2.3	3.2	.....	3.3
	121,000	8.5	20/ 8/49	15.45	16.52	13.4	8.56	3.0	3.3	.....	3.0
Woodward's Landing..... (main river channel)..	27,100	18.5	13/12/48	11.40	14.28	12.4	9.15	9.3	45.5	.....	72.8
	128,000	15.0	26/ 4/49	13.52	17.33	12.2	11.15	5.0	0.61	.....	0.49
	318,000	25.0	17/ 5/49	18.45	23.19	14.7	15.22	2.3	0.61	.....	0.73
	121,000	21.4	20/ 8/49	15.00	16.52	13.4	8.56	3.0	0.36	.....	0.36
At Fraser Ave. Bridge (Vancouver)... North arm of river	27,100	15.0	13/12/48	10.35	14.28	12.4	9.15	9.3	15.2	.....	17.5
	128,000	12.0	26/ 4/49	14.17	17.33	12.2	11.15	5.0	0.36	.....	0.30
	318,000	10.5	17/ 5/49	18.30	23.19	14.7	15.22	2.3	0.24	.....	0.24
	121,000	10.5	20/ 8/49	14.15	16.52	13.4	8.56	3.0	0.36	.....	0.36
New Westminster.....	27,100	18.0	13/12/48	13.52	14.28	12.4	9.15	9.3	1.8	.....	6.1
	128,000	10.5	26/ 4/49	15.32	17.33	12.2	11.15	5.0	0.36	.....	0.30
	318,000	11.5	17/ 5/49	17.15	23.19	14.7	15.22	2.3	0.49	.....	0.49
	121,000	9.6	20/ 8/49	13.00	16.52	13.4	8.56	3.0	0.43	.....	0.43
Mission City.....	318,000	19.4	17/ 5/49	15.30	23.19	14.7	15.22	2.3	0.49	.....	0.55
	121,000	15.4	20/ 8/49	11.0	16.52	13.4	8.56	3.0	0.49	.....	0.49
Chilliwack..... (at Rosedale ferry)	121,000	13.5	20/ 8/49	9.40	16.52	13.4	8.56	3.0	0.36	.....	0.43

\* Fraction of river depth (column 3) at which sample for chloride content was taken—for example, line 1, 15.02 p.p.m. Cl at  $\frac{1}{5}$  of 13.5 ft., or at 2.7 ft.  
† Above lowest of normal low tides over a period of at least 7 years; reference Vancouver Harbour, B.C.



TABLE I—Concluded  
Tidal Influence—Fraser River—Concluded  
B. RECEDING TIDE

Sampling Location	River Discharge at Hope, B.C. (second-feet)	River Depth at Sampling (feet)	Sampling		High Tide		Low Tide		Chloride Content, p.p.m.		
			Date	Average Time	Time	Height feet	Time	Height feet	1/5*	3/5*	4/5*
Mission City.....	27,100	23.0	13/12/48	16.20	14.28	12.4	22.17	1.9	0.61	.....	0.61
	128,000	22.6	26/ 4/49	17.52	17.33	12.2	23.01	8.0	0.24	.....	0.18
Chilliwack..... (at Rosedale Ferry)	74,600	6.0	26/10/48	16.30	13.42	12.2	19.39	8.2	.....	1.1	.....
	27,100	14.0	13/12/48	17.32	14.28	12.4	22.17	1.9	0.55	.....	0.55
	128,000	16.0	26/ 4/49	19.30	17.33	12.2	23.01	8.0	0.24	.....	0.24
	318,000	16.0	17/ 5/49	13.30	7.11	11.6	15.22	2.3	0.55	.....	0.85
Hope.....	27,100	20.0	13/12/48	18.35	14.28	12.4	22.17	1.9	0.06	.....	0.06
	128,000	26.0	26/ 4/49	20.52	17.33	12.2	23.01	8.0	0.36	.....	0.36
	318,000	23.8	17/ 5/49	14.30	7.11	11.6	15.22	2.3	0.85	.....	0.85
	121,000	18.0	20/ 8/49	8.00	0.04	12.9	8.56	3.0	0.36	.....	0.43

Table II tabulates in detail the results of chemical analyses carried out on surface waters collected at the locations shown in Figure 2 (in map pocket). An average analysis is determined for the sampling period at all daily and monthly stations. This average is the arithmetical mean of each major constituent over the period and is not weighted as to river flow. Per cent sodium and the saturation index have also been calculated for these average waters. The reader is referred to Water Survey Report No. 1 for the interpretation of per cent sodium, saturation index and other values reported in Tables II and III. Boron has also been determined occasionally to indicate the suitability of the waters for irrigation.

Figure 3 shows graphically the variation in total and non-carbonate hardness in the Fraser River.

The relationships between river discharge or level and mineral content or chemical quality of the Fraser River at Mission City, the Thompson River at Kamloops and the Quesnel River near Quesnel, are graphically shown in figures 4, 5 and 6 respectively.

## DISCUSSION

As in previous reports it is not proposed at this time to discuss in any detail the data reported in Table II. It will be noted however that most surface waters in this basin are soft or at the lower limit of medium hard using the following classification:

Soft water.....	Below 60 p.p.m. total hardness as CaCO <sub>3</sub> .
Medium hard water.....	61—120 “ “ “ “ “
Hard water.....	121—180 “ “ “ “ “
Very hard water.....	Greater than 180 p.p.m. total hardness as CaCO <sub>3</sub> .

In general, the rivers do not show any very marked variation in hardness or mineralization from season to season. However a large proportion of the waters are turbid and show considerable variation in turbidity.

Figure 3 shows graphically some decrease in hardness of Fraser River water as it approaches the sea, which can generally be explained by considering the character and volume of tributary waters entering the main river. Those which flow into Fraser River from the west or from the coastal range, are usually softer in character than those entering from the southern or eastern portion of the basin, even though the latter are not particularly hard. The head waters of Fraser River, which rise in the calcareous Rocky Mountains are, as expected from studies in the Columbia River system, somewhat harder in character. Certain tributary waters are also noticeably different in character, usually harder, as for instance, those from the Merritt area and those from the Cariboo Range or Lac la Hache area.

Figure 4 shows a close relationship between specific conductance and total hardness of Fraser River water at Mission City. The variation in both is quite small in comparison with the wide variation in flow which is paralleled by changes in turbidity. Figure 4 therefore indicates that Fraser River in flood carries a proportionately increased amount of silt but that this silt is relatively insoluble and does not affect to any great extent the mineralization of the water; that is, run-off water which produces flood conditions is not much different in dissolved mineral content than that found at normal river flow.



Figure 5 shows that a somewhat similar relationship exists in the Thompson River at Kamloops although the water is lower in hardness and total mineralization. The major difference is the lack of turbidity in the Thompson River. Even during flood the turbidity never increases enough to cause trouble for most industrial uses.

Figure 6 illustrates that tributary waters from the east central portion of the basin or Cariboo mountains are somewhat harder, but relatively constant except for a marked increase in turbidity during the flood period. During periods of low flow the Quesnel River increases in mineralization, especially sulphate ion content.

Since many of the tributary rivers are clear streams in comparison with the turbid and rapid Fraser River it is found that mixing of the tributary is often not complete for some distance downstream from the junction. A notable example of this is seen at Prince George, the clear Nechako River being visible (in the milky Fraser River) for many miles below Prince George.

Table I indicates that serious contamination of Fraser River water with seawater by tidal action is not found much above Steveston and here only when the tide is rising and river discharge is low.

The influence of incoming tide, shown by increasing chloride ion content, is noted in the river up to New Westminster when river discharge is low, but the effect is very small. From the combinations of sampling times, river discharges and tides shown in Table I it is seen that although tides may influence Fraser River levels for a considerable distance upstream they have little effect on water quality much above Steveston or possibly Woodward's Landing, insofar as industrial use is concerned.

As expected, tides entering the river tend to flow along the bottom of the river; this is indicated by the higher salinities at greater sampling depths.

### SUMMARY

Surface water supplies within the Fraser River basin are relatively constant in quality. While the main river itself differs from most British Columbia rivers by its constant turbidity it is not a hard water. Some of the cloudiness or turbidity in Fraser River water may also be due to the turbulence and saturation of the water with air and air bubbles, which gives the water a milky appearance. This turbulence also tends to disperse and finely divide the insoluble matter carried by the river so that a considerable portion is in colloidal suspension. Tributary streams except for those entering from calcareous mountain regions are generally very soft to soft in character.

In general, except for the need for clarification of the Fraser River and some tributary waters, surface waters in the basin are satisfactory for industrial use. There is generally an abundance of water available within the watershed for industrial use with little or no treatment required. However, satisfactory clarification of a turbid water such as Fraser River water does present a problem for certain uses.








FIGURE 3. GRAPH SHOWING CHANGE IN HARDNESS ALONG FRASER RIVER WATERSHED





Note. Each subdivision of a month represents  
a tri-monthly composite of daily samples





Note. Each subdivision of a month represents  
a tri-monthly composite of daily samples








TABLE II  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-foot)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>3</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION NO. 1: FRASER

1	Aug. 17/50...	116	Max. 7.85	Min.† 6.71	Max. 8.37	Min. 7.48	64	..... (9.9)*	..... (2.0)	7.7 (7.7)	15 (5)	20 (15)	38	32	80.2	54.6	0.074	.....	6.2	11.5
---	---------------	-----	-----------	------------	-----------	-----------	----	-----------------	----------------	--------------	-----------	------------	----	----	------	------	-------	-------	-----	------

\* Values in brackets refer to tests done in the mobile laboratory immediately after sampling.

STATION No. 2: FRASER RIVER

			Max.	Min.*	Max.	Min.*														
2	Aug. 24/49***	78	.....	.....	.....	.....	61	.....	.....	7.5 (1.7)	15	45	43	41	111	71.4	0.097	.....	12.8	16.8
3	Feb. 1-10/50.	32	5.19	1.10	.....	.....	33	.....	.....	7.9 (8.1)	7	0.9	.....	.....	151	105	0.143	.....	7.0	21.5
4	Feb. 11-20...	29	5.24	1.54	5.06	1.43	36	.....	.....	7.6	5	4	9.4	6.6	120	77.4	0.105	.....	9.6	16.2
5	Feb. 21-28**	.....	4.68	1.32	.....	.....	37	.....	.....	7.8	7	3	.....	.....	114	.....	.....	.....	.....	.....
6	Mar. 1-10...	31	5.55	2.31	.....	.....	37	.....	.....	7.9	5	15	32	29	116	75.2	0.102	.....	9.0	15.3
7	Mar. 11-20...	33	4.66	0.879	4.83	1.31	38	.....	.....	8.0	5	6	11	9.0	123	78.8	0.107	.....	5.8	17.0
8	Mar. 21-31...	37	4.77	0.782	.....	.....	40	.....	.....	7.7	10	3	.....	.....	119	89.4	0.122	.....	10.4	15.0
9	Apr. 1-10...	29	4.87	0.79	.....	.....	42	.....	.....	8.0	5	9	16	13	128	84.8	0.115	.....	10.6	17.2
10	Apr. 11-20...	30	4.75	1.73	4.84	1.51	43	.....	.....	8.0	5	15	44	40	124	75.8	0.103	.....	8.4	16.4
11	Apr. 21-30...	23	4.90	2.01	.....	.....	42	.....	.....	7.7	25	35	89	84	127	86.8	0.118	.....	9.6	16.6
12	May 1-10...	32	6.76	4.78	.....	.....	45	.....	.....	7.7	30	140	175	169	120	103	0.140	.....	17.6	16.7
13	May 11-20	36	11.57	10.56	10.16	8.99	48	.....	.....	7.7	35	210	261	252	113	84.0	0.114	.....	14.8	16.1
14	May 21-31...	25	11.97	11.38	.....	.....	49	.....	.....	7.6	25	70	104	99	108	78.6	0.107	.....	12.6	15.2
15	June 1-10...	37	14.49	14.04	.....	.....	52	.....	.....	7.8	10	120	128	120	104	147	0.200	.....	62.0	16.9
16	June 11-20...	55	21.00	19.64	19.34	18.58	56	.....	.....	7.8	40	200	267	258	101	74.6	0.101	.....	11.4	15.4
17	June 21-30...	45	22.55	22.07	.....	.....	55	.....	.....	7.7	25	145	187	181	99.0	71.8	0.098	.....	9.4	15.0
18	July 1-10...	71	19.96	19.62	.....	.....	59	.....	.....	7.8	20	85	106	101	91.5	71.4	0.097	.....	11.2	15.2
19	July 11-20...	71	16.01	15.57	16.46	16.07	60	.....	.....	7.9	10	70	8.8	6.0	93.1	69.8	0.095	.....	10.2	14.2
20	July 21-31...	101	13.67	11.17	.....	.....	62	.....	.....	7.9	10	50	80	77	92.3	61.8	0.084	.....	9.4	13.2
21	Aug. 1-10...	90	9.81	9.22	.....	.....	61	.....	.....	7.5	4	40	60	57	95.4	62.4	0.085	.....	15.2	13.5
22	Aug. 11-20...	92	7.92	6.95	8.37	7.48	64	.....	.....	7.6	5	30	47	45	100	67.0	0.091	.....	7.2	14.2

(\*) Discharge records are tentative data and subject to revision.

\* River flow at this station is affected by tide as shown by maximum and minimum gauge levels; see also Table I.

\*\* Results shown are preliminary tests carried out on composite sample by British Columbia Research Council; sample lost in transit.

\*\*\* Field sample, not included in average.

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.	
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		
RIVER AT HANEY																					
1.9	1.1	0.7	1.6	0.06	5.9	0 (0)	.....	0.05	.....	41.5 (40.3)	0 (0)	.....	5.8	2.5	36.5 (38.0)	47.4	6.0	.....	1.1	1	
† Discharge records at Mission City, B.C.																					
FROM BRIDGE AT MISSION CITY.																					
2.4	1.4	0.9	1.0	0.02	10.2	0 (0)	Trace	0.10	.....	54.3 (53.7)	0 (0)	5.6	6.2	7.3	51.8	65.7	.....	.....	.....	2	
4.5	2.9	1.0	.....	0.18	14.8	0	0.5	0.05	0	78.3	0	16.2	13.4	7.9	72.1	96.8	.....	.....	.....	3	
3.6	2.2	0.8	0.57	0.06	15.5	0	0.7	0.05	.....	63.2	0	9.4	6.5	3.4	55.2	76.7	.....	.....	.....	4	
.....																					
3.2	2.2	0.7	1.8	0.13	18.5	0	0.5	0.05	.....	56.6	0	8.6	8.6	1.8	51.4	74.1	.....	.....	.....	6	
3.6	2.6	0.9	1.1	0.06	16.6	0	0.5	0.10	0	62.0	0	6.6	7.5	6.4	57.2	79.3	.....	.....	.....	7	
4.0	2.7	1.3	.....	0.14	9.1	0	0	0.15	.....	68.3	0	8.2	6.2	0	53.9	72.1	.....	.....	.....	8	
5.0	2.7	0.9	0.93	0.11	13.0	0	0	0.05	0	65.9	0	8.6	9.2	9.5	63.5	80.5	.....	.....	.....	9	
3.1	3.0	0.9	1.8	0.08	15.8	0	0.4	0.15	.....	61.0	0	6.6	7.0	3.7	53.7	76.7	.....	.....	.....	10	
3.4	2.6	1.1	4.3	0.22	10.2	0	1.3	0.10	.....	65.9	0	8.6	7.0	1.4	55.4	74.9	.....	.....	.....	11	
3.3	2.4	1.2	10.9	0.68	12.8	0	0.6	0.10	.....	66.9	0	.....	7.8	0.5	55.3	78.4	.....	.....	.....	12	
3.1	2.2	1.0	17.8	0.27	6.1	0	0.7	.....	.....	64.4	0	.....	5.5	0.2	53.0	66.6	.....	.....	.....	13	
2.8	1.8	0.9	5.8	0.33	7.6	0	0.5	0.05	.....	59.0	0	.....	5.9	1.0	49.4	64.0	.....	.....	.....	14	
3.1	2.2	0.7	6.5	0.11	6.4	0	0.5	0.05	.....	57.8	0	.....	6.7	7.6	55.0	65.0	.....	.....	.....	15	
2.1	1.8	0.8	12.5	0.20	5.6	0	0.7	0.10	0	56.1	0	.....	6.3	1.0	47.0	60.6	.....	.....	.....	16	
2.4	1.6	0.7	10.2	0.27	5.8	0.2	0	0.10	.....	53.7	0	.....	5.7	3.3	47.3	58.1	.....	.....	.....	17	
2.1	1.8	0.8	5.2	0.26	9.2	0	0.4	0.10	.....	50.0	0	.....	8.8	5.5	46.5	63.3	.....	.....	.....	18	
2.5	1.6	0.7	3.7	0.22	6.9	0	0.4	0.05	.....	49.5	0	.....	6.2	5.1	45.7	57.0	.....	.....	.....	19	
2.1	2.0	0.7	3.3	0.07	5.8	0	0.6	.....	.....	54.9	0	.....	5.7	0	41.5	57.1	.....	.....	.....	20	
2.3	1.8	0.8	2.3	0.03	5.8	0	0.8	.....	0.01	48.8	0	.....	5.5	3.2	43.2	54.5	.....	.....	.....	21	
2.3	2.0	0.5	2.3	0.06	7.6	0	0	0.10	.....	52.7	0	.....	7.7	1.7	44.9	60.3	.....	.....	.....	22	

(b) Total refers to sum of hardness due to calcium and magnesium ions.

(c) Total iron increases with turbidity in many waters, indicating analyses of iron in colloidal and suspended matter.



TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)				Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean									Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 2: FRASER RIVER FROM

	1950		Max.	Min.	Max.	Min.														
1	Aug. 21-31...	93	7.49	6.37	.....	.....	64	.....	.....	7.8	15	20	48	47	99.8	68.6	0.093	.....	6.2	15.2
2	Sept. 1-10...	97	6.41	5.44	.....	.....	62	.....	.....	7.8	5	50	56	50	106	68.4	0.093	.....	4.8	15.8
3	Sept. 11-20...	73	5.70	3.62	5.82	4.10	60	.....	.....	7.9	5	20	32	30	106	70.8	0.096	.....	6.0	15.7
4	Sept. 21-30...	63	5.35	3.25	.....	.....	58	.....	.....	7.7	2	45	26	25	103	66.4	0.090	.....	7.0	15.0
5	Oct. 1-10...	67	5.42	2.93	.....	.....	52	.....	.....	7.7	2	35	34	30	107	68.4	0.093	.....	6.6	15.1
6	Oct. 11-20...	110	6.41	4.30	5.96	3.58	49	.....	.....	7.6	20	20	33	30	99.7	65.4	0.089	.....	9.8	14.2
7	Oct. 21-31...	100	6.03	3.51	.....	.....	46	.....	.....	7.6	10	9	18	16	110	78.0	0.106	.....	11.2	15.2
8	Nov. 1-10...	109	6.32	4.05	.....	.....	43	.....	.....	7.7	15	50	67	61	104	76.8	0.104	.....	16.6	18.0
9	Nov. 11-20...	89	6.38	3.23	6.36	3.65	39	.....	.....	7.6	5	25	36	34	108	70.6	0.096	.....	9.4	16.0
10	Nov. 21-30...	73	6.37	3.65	.....	.....	40	.....	.....	7.4	15	15	29	27	93.3	61.8	0.084	.....	11.6	13.4
11	Dec. 1-10...	53	6.45	3.50	.....	.....	37	.....	.....	7.5	5	6	14	12	103	66.4	0.090	.....	11.0	14.0
12	Dec. 11-20...	59	6.94	4.26	7.13	4.51	40	.....	.....	7.3	10	20	13	11	90.3	61.0	0.083	.....	15.2	12.9
13	Dec. 21-31...	43	7.92	5.65	.....	.....	42	.....	.....	7.6	15	10	34	30	82.3	57.4	0.078	.....	10.8	11.7
14	Jan. 1-10/51	52	5.98	2.72	.....	.....	37	.....	.....	7.5	10	7	12	11	101	66.8	0.091	.....	10.8	13.8
15	Jan. 11-20...	42	5.74	1.81	5.81	2.15	37	.....	.....	7.4	10	3	.....	.....	116	81.2	0.111	.....	12.8	15.6
16	Jan. 21-31...	43	5.70	1.95	.....	.....	36	.....	.....	7.6	10	9	19	15	119	67.6	0.092	.....	9.6	14.7
17	Yearly Average (35 Samples)	59	8.36	6.15	8.34	6.11	47	.....	.....	7.7	10	45	.....	.....	107.9	76.0	0.103	.....	11.7	15.3

STATION No. 3: FRASER RIVER

			Gauge height in feet																	
18	Aug. 26/49*	76	.....	.....	62	.....	.....	7.5	15	50	50	49	109	73.0	0.099	.....	12.0	19.6		
19	Apr. 5/50†	12	1.57	2.64	39	.....	(1.5)	(8.3)	7	(45)	18	18	153	93.2	0.127	.....	7.2	20.5		
20	Apr. 24.....	15	3.32	2.64	40	.....	.....	8.0	20	200	159	153	146	97.2	0.132	.....	8.6	20.3		
21	May 29.....	38	13.24	11.56	48	.....	.....	7.9	10	70	.....	.....	116	.....	.....	.....	.....	15.8		
22	June 23.....	42	19.42	18.48	57	.....	.....	7.8	40	150	163	158	98.9	75.0	0.102	.....	9.2	15.5		
23	July 24.....	81	14.18	15.37	61	.....	.....	7.7	5	105	97	92	98.8	65.4	0.089	.....	9.8	16.9		

\* Field sample, not included in average.

† Extra sample, not included in average.

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-	
2.4	1.8	0.7	2.0	0.05	7.6	0	0.4	0.10	.....	53.7	0	.....	6.8	3.8	47.8	61.4	.....	.....	.....	1
2.6	1.8	1.0	3.2	0.04	9.4	0	0.5	0	.....	53.7	0	.....	.....	6.1	50.1	57.5	.....	.....	.....	2
2.7	1.6	0.7	1.3	0.06	8.6	0	0.4	0.05	.....	59.0	0	.....	7.0	1.9	50.3	65.8	.....	.....	.....	3
2.6	1.8	0.7	1.0	0.06	8.6	0	0.4	0.05	.....	55.6	0	.....	4.8	2.5	48.1	62.5	.....	.....	.....	4
2.8	1.7	0.9	2.2	0.05	8.4	0	0	0.10	.....	56.1	0	.....	5.1	3.2	49.2	64.0	.....	.....	.....	5
2.8	2.2	1.0	1.5	0.10	8.2	0	0.5	0.10	.....	53.2	0	.....	7.0	3.4	47.0	63.8	.....	.....	.....	6
2.9	2.0	0.6	0.8	0.05	8.2	0	0.5	0.10	.....	57.3	0	.....	5.7	3.0	50.0	64.3	.....	.....	.....	7
2.9	2.3	0.8	1.4	0.09	17.6	0	0	.....	0	56.1	0	.....	3.0	10.8	56.8	73.2	.....	.....	.....	8
2.9	1.3	0.4	1.8	0.09	9.5	0	0.6	.....	.....	58.8	0	.....	7.5	3.6	51.8	69.0	.....	.....	.....	9
2.6	2.1	0.5	2.2	0.09	6.7	0	0.9	0.05	.....	51.2	0	.....	5.8	2.1	44.1	59.6	.....	.....	.....	10
2.8	2.3	0.3	0.9	0.10	6.3	0	0.9	0.05	.....	55.6	0	.....	6.7	0.8	46.4	61.8	.....	.....	.....	11
2.5	3.3	0.5	0.7	0.05	9.1	0	0.6	0.10	.....	48.8	0	.....	6.7	2.5	42.5	60.5	.....	.....	.....	12
2.3	1.8	0.5	1.7	0.11	4.4	0	1.3	0.05	.....	43.7	0	.....	5.2	2.9	38.7	50.6	.....	.....	.....	13
2.5	1.5	0.6	0.99	0.03	7.2	0	0.4	0.10	.....	51.5	0	4.4	4.6	2.5	44.7	61.5	.....	.....	.....	14
3.1	1.9	0.6	.....	0.28	8.0	0	0.5	0.05	.....	60.0	0	.....	4.4	2.5	51.7	64.1	.....	.....	.....	15
3.1	1.9	0.5	1.2	0.06	10.9	0	0.7	0	.....	54.9	0	7.8	7.2	4.4	49.4	66.0	.....	.....	.....	16
2.9	2.1	0.8	.....	0.13	9.5	0	0.5	0.08	.....	57.3	0	.....	6.6	3.1	50.1	66.0	8.2	.....	0.8	17

## FROM FERRY AT ROSEDALE

2.9	1.8	0.9	0.75	0.08	15.7	0	Trace	0.10	.....	59.5 (61.0)	0 (0)	5.0	6.2	12.0	60.8	76.5	.....	.....	.....	18
4.9	2.8	1.0	2.2	0.06	17.4	0	0.4	0.10	.....	80.8	0	5.8	6.6	5.1	71.3	93.5	.....	.....	.....	19
4.0	3.6	1.2	8.8	0.23	10.7	0	0.5	0.10	.....	78.1	0	9.2	8.5	3.1	67.1	87.5	.....	.....	.....	20
2.8	2.6	1.0	.....	.....	7.6	0	.....	.....	0	65.4	0	.....	7.2	0	50.9	69.2	.....	.....	.....	21
2.3	1.4	0.7	4.6	0.31	6.4	0	0.4	0.05	.....	56.1	0	.....	6.0	2.2	48.2	55.2	.....	.....	.....	22
2.2	1.8	0.7	1.2	0.14	6.4	0	0.4	.....	.....	56.6	0	.....	5.2	4.9	51.3	61.5	.....	.....	.....	23



TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water tem- pera- ture (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conduct- ance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre- foot	Thou- sand tons per day		

STATION No. 3: FRASER RIVER

			Gauge height in feet															
1	Aug. 17*	111	9.32	9.79	65			7.4 (2.0)	15	40 (40)			109					15.9
2	Aug. 25	71	8.53	9.79	63			7.7 (8.0)	10	50			111					16.7
3	Sept. 25	70	5.80	6.73	59			8.0	5	25			114					16.3
4	Oct. 23	91	4.62	5.35	45			7.8	5	20	20	19	123	78.4	0.107		8.8	17.9
5	Nov. 25	65	3.56	5.16	32			7.5	15	15			116					17.5
6	Dec. 23/50	46	4.70	4.51	41			7.6	15	15			108					15.8
7	Feb. 5/51	8	2.44	2.74	34			7.5	5	0.2			138					22.0
8	Feb. 23	15	2.16	2.74	35			8.5	10	9			156					20.5
9	Mar. 24	114	2.06	1.93	38			7.9	10	560	341	319	180	114	0.155		34.4	23.6
10	Yearly Average (12 samples)	55	7.01	7.25	46			7.8	13	100			125.5	86.0	0.117		14.2	18.2

\* Field sample, not included in average.

STATION No. 4: FRASER RIVER FROM BRIDGE

11	Mar. 8/50*	8	23,100	23,600	41			7.8	7	15	32	30	141	88.6	0.121	5.54	9.2	19.0
12	Mar. 23	3	23,900	23,600	39			8.2	5	15	91	39	156	96.2	0.131	6.20	10.0	20.4
13	Apr. 22	17	41,800	33,600	40			8.0	20	310	196	190	141	94.8	0.129	10.7	9.0	19.5
14	May 23	9	160,000	142,000	40			7.6	20	115			94.9					15.0
15	June 23	34	427,000	309,000	48			7.9	20	140	144	137	75.5	55.8	0.076	64.3	6.4	12.4
16	July 22	83	197,000	231,000	53			7.7	2	85	97	92	95.5	64.0	0.087	34.0	21.0	15.0
17	Aug. 10**	118	121,000	115,000	63			8.0 (1.5)	8	45 (60)			103					16.0
18	Aug. 18	62	105,000	115,000	55			7.5	7	30			98.5					15.0
19	Sept. 23	66	58,200	73,900	61			7.9	5	25			116					16.1
20	Oct. 23	91	43,500	55,300	46			7.8	5	25	24	23	113	74.8	0.102	9.79	10.6	16.2
21	Nov. 24	66	35,900	53,800	40			7.5	20	20			103					15.3
22	Dec. 22	47	43,100	45,300	42			7.5	20	20			102					15.3
23	Jan. 1951	No sample taken		30,100														
24	Feb. 24	12	25,500	32,900	36			7.1	5	6	3.0	2.7	138	93.6	0.127	6.44	14.8	20.0
25	Mar. 12	29			38			8.0	5	4			138					19.9
26	Yearly Average (12 samples)	43	106,445	95,458	45			7.7	11	66			114	79.9	0.109		12.0	16.7

\* Extra sample, not included in average; low water sample.

\*\* Field sample, not included in average.

TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

Magnesium (Mg)	Alkalia		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total <sup>b</sup>			+	-		

FROM FERRY AT ROSEDALE—Continued

2.7	2.0	0.7	.....	.....	8.6	0	.....	.....	.....	59.5 (54.9)	0 (0)	.....	6.0	2.0 (7.0)	50.8 (52.0)	65.2	.....	.....	.....	.....	1
2.7	1.8	0.6	.....	.....	8.8	0	.....	.....	0.01	57.8	0	.....	4.8	5.4	52.8	63.8	.....	.....	.....	.....	2
3.1	1.8	0.6	.....	.....	8.0	0	.....	.....	.....	61.5	0	.....	5.5	3.1	53.5	65.5	.....	.....	.....	.....	3
3.7	2.0	0.8	1.3	0.06	8.9	0	0.4	0.05	.....	65.9	0	.....	6.4	5.9	59.9	72.6	.....	.....	.....	.....	4
3.8	2.2	0.9	.....	.....	10.1	0	.....	.....	0	75.6	0	.....	5.6	0	59.5	77.3	.....	.....	.....	.....	5
3.2	1.7	0.4	.....	.....	9.5	0	.....	.....	.....	58.6	0	.....	5.1	4.5	52.5	64.5	.....	.....	.....	.....	6
4.6	2.0	0.8	.....	.....	11.8	0	.....	.....	.....	83.0	0	.....	6.0	5.8	73.8	88.0	.....	.....	.....	.....	7
4.6	2.1	0.7	.....	.....	14.4	0.6	.....	.....	.....	79.3	0	.....	7.2	5.1	70.1	89.1	.....	.....	.....	.....	8
6.3	4.1	1.3	18.8	0.18	17.1	0.6	0.7	0	.....	90.0	0	.....	10.3	11.0	84.8	108	.....	.....	.....	.....	9
3.6	2.3	0.8	.....	0.08	10.0	0.1	0.5	0.04	.....	69.0	0	.....	6.6	3.7	60.3	76.2	7.5	.....	.....	0.6	10

AT HOPE—Drainage area, 85,600 square miles

4.6	2.6	0.8	1.5	0.11	14.0	0	0.5	0.10	.....	73.7	0	8.2	6.8	5.9	66.3	87.0	.....	.....	.....	.....	11
5.1	3.1	0.8	2.4	0.04	18.9	0	0.4	0.10	.....	73.7	2.9	8.6	7.1	6.7	71.9	95.1	.....	.....	.....	.....	12
3.8	3.7	1.3	11.1	0.21	11.0	0	0.4	0.10	.....	74.4	0	8.0	8.2	3.3	64.3	84.8	.....	.....	.....	.....	13
2.7	2.0	0.8	.....	.....	8.1	0	.....	.....	.....	56.6	0	.....	6.4	2.1	48.5	62.8	.....	.....	.....	.....	14
1.2	1.4	0.6	7.4	0.18	6.6	0	0.4	0.10	.....	41.5	.....	.....	5.1	1.8	35.8	48.3	.....	.....	.....	.....	15
2.3	1.6	0.8	3.0	0.12	5.8	0	0.4	.....	.....	53.7	0	.....	5.6	2.9	46.9	58.0	.....	.....	.....	.....	16
2.4	1.7	0.7	.....	.....	7.6	0	.....	.....	.....	53.7	0	.....	5.5	5.8	49.8	60.3	.....	.....	.....	.....	17
2.3	1.5	0.6	.....	.....	8.0	0	.....	.....	0	51.0	0	.....	6.7	5.1	46.9	59.2	.....	.....	.....	.....	18
3.3	2.0	0.7	.....	.....	10.5	0	.....	.....	.....	61.0	0	.....	5.2	3.8	53.8	67.8	.....	.....	.....	.....	19
3.3	1.8	0.7	1.3	0.08	8.4	0	0.4	0	.....	61.0	0	.....	6.8	4.0	54.0	67.6	.....	.....	.....	.....	20
3.1	2.0	0.8	.....	.....	7.0	0	.....	.....	0.01	63.4	0	.....	4.6	0	51.0	64.0	.....	.....	.....	.....	21
2.9	1.7	0.3	.....	.....	8.7	0	.....	.....	.....	53.7	0	.....	5.2	6.0	50.0	60.5	.....	.....	.....	.....	22
.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	23
4.0	2.3	0.8	0.59	0.05	9.9	1.5	0.7	0	0	70.8	0	.....	9.1	8.4	66.4	83.1	.....	.....	.....	.....	24
4.2	2.0	1.0	.....	.....	6.2	1.0	1.3	.....	.....	75.2	0	.....	17.2	5.3	66.9	89.8	.....	.....	.....	.....	25
3.2	2.1	0.8	.....	0.11	9.1	0.2	0.6	0.06	.....	61.3	0	.....	7.3	4.6	54.8	70.4	7.6	.....	.....	0.7	26



TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
STATION No. 5: FRASER RIVER																		
1	Aug. 10/50...	98	.....	.....	62	.....	..... (1.0)	8.0 (8.0)	5	45 (70)	56	53	105	67.0	0.091	.....	7.6	15.0
STATION No. 6: FRASER																		
2	Aug. 27/49...	75	.....	.....	61	.....	..... (2.0)	7.6 (8.3)	20	50 (20)	79	75	110	75.8	0.103	.....	15.2	17.2
STATION No. 7: FRASER RIVER																		
			Gauge height in feet															
3	Feb. 21/50*	16	.....	.....	32	.....	.....	7.4	0	0.4	.....	.....	33.6	20.4	0.028	.....	1.0	4.9
4	Mar. 21.....	6	.....	.....	38	.....	.....	7.9	5	5	8.6	7.2	289	180	0.245	.....	16.8	36.7
5	Apr. 21.....	18	.....	.....	35.5	.....	.....	7.9	25	290	261	252	169	114	0.155	.....	10.4	22.9
6	May 23.....	7	19.26	18.23	47	.....	.....	8.1	40	165	.....	.....	123	.....	.....	.....	18.0	.....
7	June 21.....	36	31.52	26.52	56.5	.....	.....	8.1	15	335	376	358	108	74.0	0.099	.....	8.4	17.4
8	July 24.....	81	20.75	.....	62	.....	.....	7.7	10	145	131	124	109	67.6	0.092	.....	12.2	16.9
9	Aug. 18.....	110	14.94	.....	64	.....	..... (1.5)	7.8 (8.2)	3	50 (40)	.....	.....	124	.....	.....	.....	18.8	.....
10	Sept. 21.....	68	10.26	.....	59	.....	.....	8.0	5	35	.....	.....	126	.....	.....	.....	18.1	.....
11	Oct. 23.....	91	8.27	.....	40	.....	.....	7.8	15	10	22	21	134	85.4	0.116	.....	9.8	20.0
12	Nov. 21.....	69	4.35	.....	34	.....	.....	7.8	15	15	.....	.....	151	.....	.....	.....	22.9	.....
13	Dec. 23.....	46	.....	.....	33	.....	.....	7.7	15	20	.....	.....	158	.....	.....	.....	22.5	.....
14	Jan. ....	No sample taken.		.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
15	Feb. 21/51...	27	.....	.....	35	.....	.....	7.4	10	7	1.8	1.2	174	109	0.148	.....	12.8	24.4
16	Yearly Average (11 samples)	51	.....	.....	46	.....	.....	7.8	15	100	.....	.....	151	.....	0.143	.....	.....	21.7
* Not included in average. Sampled through hole in ice with probable dilution with melted snow and ice.																		
STATION No. 8: FRASER RIVER AT																		
17	Aug. 8/50...	113	.....	.....	64	.....	..... (1.5)	7.7 (8.1)	6 (25)	55 (40)	53	46	120	77.4	0.105	.....	5.8	19.2
STATION No. 9: FRASER RIVER																		
18	Aug. 21/50...	134	52,200†	61,100†	64	..... (9.2)	..... (1.2)	8.0 (8.0)	5	45 (30)	43	39	122	75.4	0.103	10.60	27.4	18.0

TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total <sup>b</sup>			+		-	
FROM BRIDGE AT SPUZZUM																					
2.7	1.6	0.5	2.9	0.10	8.2	0	0.4	0.05	.....	56.1	0	.....	5.7	2.5 (4.0)	48.5 (46.0)	61.8	6.5	.....	0.5	1	
RIVER AT BOSTON BAR																					
3.0	1.8	1.0	1.3	0.13	11.2	0	0.5	0.05	.....	59.1 (61.0)	0 (0)	6.2	6.0	6.8 (1.3)	55.2 (51.3)	70.0	6.5	.....	0.95	2	
FROM BRIDGE AT LILLOOET																					
0.9	0.5	0.1	.....	0.04	4.1	0	0	Trace	Trace	17.1	0	2.4	2.0	1.9	15.9	20.9	.....	.....	.....	3	
13.0	5.6	1.2	0.57	0.10	37.0	0	2.7	0.05	.....	139	0	7.6	6.5	30.9	145	171	.....	.....	.....	4	
5.0	4.3	1.2	14.4	0.20	10.5	0	0.4	0.1	.....	94.2	0	10.8	10.2	0.5	77.7	101	.....	.....	.....	5	
3.6	1.8	0.9	.....	.....	8.4	0	.....	.....	0.03	66.9	2.4	.....	6.1	0.9	59.7	74.1	.....	.....	.....	6	
3.2	1.4	0.9	21.0	0.21	9.2	0	0.4	0	.....	56.6	2.4	.....	5.2	6.2	56.6	68.1	.....	.....	.....	7	
2.9	1.8	0.6	1.8	0.12	6.3	0	Trace	.....	.....	59.0	0	.....	5.1	5.7	54.1	62.7	.....	.....	.....	8	
3.3	1.7	0.6	.....	.....	9.1	0 (0)	.....	.....	.....	67.3 (63.4)	0 (0)	.....	4.5	5.3	60.5	71.1	.....	.....	.....	9	
3.9	2.0	0.5	.....	.....	11.9	0	.....	.....	0.005	70.5	0	.....	5.2	3.4	61.2	76.3	.....	.....	.....	10	
4.4	2.0	0.6	1.2	0.04	9.4	0	0	0.1	.....	78.1	0	.....	6.4	4.0	68.0	81.3	.....	.....	.....	11	
4.9	2.4	0.7	.....	.....	10.3	0	.....	.....	0.02	85.9	0	.....	5.6	7.1	77.5	88.5	.....	.....	.....	12	
5.2	2.0	0.5	.....	.....	10.5	0	.....	.....	.....	88.1	0	.....	5.8	5.3	77.5	89.8	.....	.....	.....	13	
																				14	
6.3	2.6	0.6	0.52	0.08	14.0	0.5	0.7	0	.....	95.2	0	9.0	7.1	8.8	86.8	103	.....	.....	.....	15	
5.1	2.5	0.75	.....	0.11	12.4	0	.....	.....	.....	82.8	0	.....	5.2	7.3	75.1	89.7	6.7	.....	0.4	16	
BRIDGE NEAR WILLIAMS LAKE																					
3.1	1.1	0.6	2.0	0.07	8.9	0 (0)	0.4	0.05	.....	67.1 (63.4)	0 (0)	.....	5.1	5.7 (7.0)	60.7 (59.0)	71.5	3.7	.....	0.7	17	
FROM FERRY AT MACALISTER																					
3.0	1.8	0.5	2.2	0.05	8.6	0	0.7	.....	.....	64.4 (59.8)	0 (0)	.....	5.2	4.4 (11.0)	57.2 (60.0)	69.5	6.3	.....	0.5	18	



TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 10: FRASER RIVER AT BRIDGE

			Gauge height in feet															
1	Mar. 19/50...	13	Ice conditions		33	.....	.....	7.9	2	5	15	12	153	90.4	0.123	.....	6.4	25.4
2	Apr. 19.....	15	5.48	5.90	44	.....	.....	8.1	15	155	214	201	154	102	0.139	.....	17.8	24.0
3	May 21.....	22	13.63	13.44	52	.....	.....	7.7	20	130	125	116	125	90.6	0.123	.....	39.4	18.9
4	June 19.....	51	21.20	17.78	64	.....	.....	7.7	5	180	237	231	115	83.6	0.114	.....	11.2	19.2
5	July 23.....	82	13.89	14.66	.....	.....	.....	7.8	5	75	98	93	112	71.2	0.097	.....	7.8	20.0
6	Aug. 22.....	115	9.19	10.24	66	.....	.....	7.6	7	50	.....	.....	120	.....	.....	.....	.....	17.7
7	Sept. 5.....	84	9.84	8.03	58	.....	(2.0)	(8.2)	10	(25)	135	.....	125	.....	.....	.....	.....	20.2
8	Sept. 23.....	66	6.95	8.03	55	.....	.....	8.0	7	35	.....	.....	129	.....	.....	.....	.....	19.2
9	Oct. 28.....	97	6.34	6.80	39	.....	.....	7.9	15	15	27	24	137	96.6	0.131	.....	11.2	21.0
10	Nov.—Sample lost in transit.				6.44	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
11	Dec. 21.....	43	5.69	5.69	36	.....	.....	7.5	10	15	.....	.....	142	.....	.....	.....	.....	23.3
12	Jan. 51—No sample taken; river frozen over.				.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
13	Feb. 19.....	29	Ice conditions		34	.....	.....	7.5	15	5	7.6	6.2	163	102	0.139	.....	12.0	23.8
14	Average..... (11 samples)	56	10.2	10.1	48	.....	.....	7.8	20	70	.....	.....	134	90.9	0.124	.....	15.1	21.2

\* Above inflow of Quesnel River.

STATION No. 11: FRASER RIVER BELOW MOUTH

			Gauge height in feet															
15	Feb. 17/50...	15	Ice conditions		33	.....	.....	8.1	8	1	.....	.....	187	123	0.167	.....	13.0	29.4
16	Mar. 16.....	11	17.01	16.79	34	.....	.....	8.0	10	3	.....	.....	199	123	0.167	.....	8.2	30.0
17	Apr. 17.....	15	14.30	14.65	33	.....	.....	8.0	0	0.5	.....	.....	131	83.8	0.114	.....	5.4	19.8
18	May 16.....	14	25.70	22.79	44	.....	.....	7.4	40	45	.....	.....	108	.....	.....	.....	.....	17.0
19	June 18.....	39	31.79	27.93	51	.....	.....	7.8	25	170	152	143	115	72.2	0.098	.....	6.8	18.9
20	July 16.....	57	24.45	24.74	54	.....	.....	7.8	10	20	.....	.....	125	78.0	0.106	.....	7.0	19.4
21	Aug. 18.....	78	19.45	19.78	55	.....	.....	7.8	15	50	.....	.....	137	.....	.....	.....	.....	22.1
22	Average..... (7 samples)	33	22.1	21.1	43	.....	.....	7.8	15	75	.....	.....	143	96.0	0.130	.....	8.1	22.4

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Totale	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>						
																		+	-		

AT QUESNEL\*—Drainage area, 38,000 square miles

3.9	1.6	0.2	0.40	0.04	15.2	0	0.7	0	.....	84.2	0	4.4	4.6	10.5	79.5	93.1	.....	.....	.....	.....	1
5.0	2.0	0.9	30.0	0.25	11.9	0	0	0.30	.....	87.8	0	8.0	6.4	8.5	80.5	94.0	.....	.....	.....	.....	2
2.9	1.6	0.8	1.3	0.35	3.5	0	0.9	0.20	0	68.6	0	6.6	3.9	2.9	59.1	70.8	.....	.....	.....	.....	3
2.5	1.0	0.5	12.6	0.45	5.8	0	0.4	0	.....	63.4	0	.....	4.8	6.2	58.2	65.9	.....	.....	.....	.....	4
2.8	1.6	0.5	1.6	0.09	7.1	0	0.4	.....	.....	62.2	0	.....	3.5	10.4	61.4	66.6	.....	.....	.....	.....	5
3.1	1.5	0.5	.....	.....	8.8	0	.....	.....	.....	64.7 (63.4)	0 (0)	.....	4.1	4.0	57.0	68.0	.....	.....	.....	.....	6
3.5	1.2	0.5	.....	.....	9.5	0	.....	.....	0.007	69.8	0	.....	4.6	7.6	64.8	74.5	.....	.....	.....	.....	7
3.6	2.0	0.7	.....	.....	8.8	0	.....	.....	.....	73.2	0	.....	5.3	2.7	62.7	76.0	.....	.....	.....	.....	8
4.5	2.2	0.3	1.1	0.07	9.1	0	0.5	0.10	.....	82.5	0	.....	10.1	3.4	71.0	88.5	.....	.....	.....	.....	9
.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	10
4.6	1.9	0.6	.....	.....	14.8	0	.....	.....	.....	81.5	0	.....	4.6	10.2	77.0	90.3	.....	.....	.....	.....	11
.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	12
5.3	2.1	0.5	0.79	0.10	12.5	1.0	0.7	0	.....	95.2	0	10.0	7.2	3.2	81.2	100	.....	.....	.....	.....	13
3.8	1.7	0.6	.....	0.19	9.7	0	0.5	0.10	.....	75.7	0	.....	5.4	6.5	68.5	80.7	5.1	.....	.....	0.5	14

OF NECHAKO RIVER AT PRINCE GEORGE

5.6	2.2	0.6	.....	0.10	12.4	0	0.3	0	0.08	104	2.4	6.0	6.2	7.6	96.4	110	.....	.....	.....	.....	15
6.5	2.4	0.6	.....	0.32	17.3	0	0.5	0.05	.....	114	0	8.4	7.7	7.8	102	121	.....	.....	.....	.....	16
3.1	2.7	0.3	.....	0.09	13.5	0	0.6	0.10	.....	66.6	0	7.8	5.9	7.6	62.2	78.8	.....	.....	.....	.....	17
2.7	1.1	0.6	.....	.....	9.3	0	.....	.....	0.01	63.4	0	.....	4.4	1.5	53.5	66.3	.....	.....	.....	.....	18
2.7	0.7	0.6	6.8	0.18	6.4	0	0.5	0	.....	64.9	0	.....	3.5	5.1	58.3	65.3	.....	.....	.....	.....	19
3.7	1.4	0.7	1.5	0.41	9.9	0.5	1.3	0.05	.....	67.8	0	5.2	2.7	8.0	63.6	71.9	.....	.....	.....	.....	20
3.4	1.0	0.3	.....	.....	9.1	0	.....	.....	0.02	73.0	0	.....	4.9	9.3	69.1	76.7	.....	.....	.....	.....	21
4.0	1.6	0.5	.....	0.22	11.1	.....	.....	.....	.....	83.4	0.3	.....	5.0	4.0	72.3	84.3	4.5	.....	.....	0.4	22



TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K × 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 12: FRASER RIVER ABOVE

			Gauge height in feet															
1	Feb. 17/50...	15	.....	.....	33	.....	.....	7.7	5	3	.....	.....	214	133	0.181	.....	13.4	35.8
2	Mar. 16.....	11	.....	.....	33	.....	.....	8.1	5	0.4	.....	.....	339	209	0.284	.....	15.4	46.1
3	Apr.—No sample taken.		.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
4	May 16.....	14	78,000†	57,300†	44	.....	.....	7.5	45	45	.....	.....	109	.....	.....	.....	.....	17.4
5	June—No sample taken.		.....	92,400	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
6	July 16.....	66	54,660	57,400	55	.....	.....	7.8	15	25	66	63	122	72.4	0.098	10.65	5.0	19.5
7	Aug.—No sample taken.		.....	31,300	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
8	Sept. 2.....	131	29,800	21,300	55	.....	.....	7.9 (2.0)	15	50 (7.7)	.....	.....	138	.....	.....	.....	.....	21.4
9	Sept. 15.....	70	16,400	21,300	38	.....	.....	7.9	10	25	.....	.....	153	.....	.....	.....	.....	24.0
10	Oct.—Sample lost in transit.		.....	15,700	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
11	Nov.—No sample taken.		.....	18,600	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
12	Dec. 12.....	45	8,700	8,020	35	.....	.....	7.7	10	15	.....	.....	184	.....	.....	.....	.....	28.3
13	Jan. 9/51...	43	6,200	5,610	34	.....	.....	7.7	15	3	.....	.....	193	117	0.159	1.96	16.6	30.2
14	Feb. 10.....	38	4,900	4,820	38	.....	.....	7.8	7	0.9	.....	.....	213	127	0.173	1.68	9.2	33.8
15	Average..... (9 samples)	48	28,370	34,770	41	.....	.....	7.8	15	20	.....	.....	185	132	0.179	.....	11.9	28.5

† Discharge records at Shelley, drainage area 12,500 square miles; station established May 2, 1950.

STATION No. 13: FRASER RIVER AT

16	Feb. 16/50*..	16	Low†	.....	33	.....	.....	8.2	5	4	.....	.....	195	.....	.....	.....	.....	.....
17	Mar. 16.....	.....	Low	.....	34	.....	.....	8.0	5	3	.....	.....	194	116	0.158	.....	7.0	26.0
18	Apr. 15.....	12	Low	.....	38	.....	.....	7.8	5	5	4.6	3.2	195	118	0.161	.....	10.4	28.0
19	May 16.....	27	Normal	.....	42	.....	.....	7.8	7	7	19	13	152	89.8	0.121	.....	33.6	20.0
20	June 15.....	27	High	.....	50	.....	.....	8.0	10	120	110	104	131	242	0.329	.....	39.6	34.2
21	July 15.....	58	High	.....	47	.....	.....	8.1	25	35	74	72	113	77.2	0.105	.....	5.2	17.0
22	Aug. 15.....	65	Normal	.....	52	.....	.....	8.1	10	70	.....	.....	112	.....	.....	.....	.....	17.8
23	Sept. 16.....	69	Normal	.....	51	.....	.....	8.0	15	40	.....	.....	121	.....	.....	.....	.....	18.3
24	Oct. 14.....	52	Low	.....	44	.....	.....	7.8	2	0	10.4	9.0	134	82.2	0.112	.....	16.4	17.6

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts<sub>per</sub> million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		

## MOUTH OF NECHAKO RIVER AT PRINCE GEORGE

6.5	2.2	0.7	.....	0.15	15.0	0	0.8	0	0.13	123	0	6.8	6.8	15.6	116	128	.....	.....	.....	.....	1
15.5	5.1	2.3	.....	0.09	17.6	5.1	0.4	0	.....	198	4.8	16.6	14.9	8.8	179	209	.....	.....	.....	.....	2
																					3
2.7	0.9	0.6	.....	.....	8.4	0	.....	.....	0.21	62.5	0	.....	4.3	3.3	54.5	65.0	.....	.....	.....	.....	4
																					5
3.2	0.8	0.3	4.9	0.18	9.4	0	0.4	0	.....	65.9	0	.....	5.0	7.9	61.9	71.1	.....	.....	.....	.....	6
																					7
4.6	1.1	0.8	.....	.....	9.1	0	0	.....	.....	75.6 (75.6)	0 (0)	.....	3.8	10.3	72.3	78.0	.....	.....	.....	.....	8
4.1	1.4	0.4	.....	.....	10.1	0	.....	.....	.....	90.3	0	.....	5.0	2.8	76.8	89.4	.....	.....	.....	.....	9
																					10
																					11
5.7	1.7	0.6	.....	.....	10.3	0	.....	.....	.....	104	0	.....	5.0	8.4	94.0	103	.....	.....	.....	.....	12
5.6	1.8	0.4	.....	0.09	8.6	0	0	.....	.....	111	0	.....	1.9	7.2	98.4	103	.....	.....	.....	.....	13
6.9	1.9	0.4	.....	0.09	13.0	0	1.1	0	.....	125	0	6.8	6.5	10.3	113	140	.....	.....	.....	.....	14
6.1	1.9	0.7	.....	0.12	11.3	.....	0.45	0	.....	106	0.5	.....	5.9	9.3	96.2	110	4.1	.....	.....	0.2	15

## BRIDGE NEAR TÊTE JAUNE CACHE

.....	.....	.....	.....	.....	30.9	0	.....	.....	.....	97.4	2.4	.....	.....	17.6	101	.....	.....	.....	.....	.....	16
8.0	2.0	0.9	.....	0.20	21.2	0	0.4	0	0	97.1	0	6.0	4.7	18.2	97.8	111	.....	.....	.....	.....	17
7.3	2.0	1.2	0.47	0.06	30.8	0	0.4	0.10	.....	100	0	5.0	4.6	17.9	99.9	124	.....	.....	.....	.....	18
5.2	1.2	0.9	0.30	0.07	15.8	0	0.8	0.05	0.03	73.2	0	4.0	2.9	11.3	71.3	83.4	.....	.....	.....	.....	19
5.9	1.1	1.1	10.4	0.08	12.5	0	0.7	0	.....	68.6	0	.....	3.2	53.4	110	92.4	.....	.....	.....	.....	20
4.2	0.7	0.8	1.6	0.42	11.0	0	0.4	0	.....	55.4	0	8.0	2.4	14.3	59.7	64.1	.....	.....	.....	.....	21
3.0	0.8	0.9	.....	.....	12.1	0	.....	.....	0.02	58.6	0	.....	3.6	8.7	56.7	67.0	.....	.....	.....	.....	22
4.0	0.7	0.6	.....	.....	12.6	0	.....	.....	.....	65.9	0	.....	2.9	7.8	62.2	71.5	.....	.....	.....	.....	23
4.5	0.8	0.6	0.68	0.04	15.0	0	0.4	0.16	.....	67.1	0	.....	2.9	7.4	62.4	74.9	.....	.....	.....	.....	24



TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-foot)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
STATION No. 13: FRASER RIVER AT																		
1	Nov. 14/50...	76	Low	.....	32	.....	.....	7.7	4	6	.....	.....	151	.....	.....	.....	.....	20.5
2	Dec. 14.....	50	Low	.....	33	.....	.....	7.6	5	8	.....	.....	163	.....	.....	.....	.....	21.3
3	Jan. 16/51...	28	Low	.....	32	.....	.....	7.7	7	1	.....	.....	176	111	0.151	.....	12.4	24.6
4	Average..... (11 samples)	43	.....	.....	41	.....	.....	7.9	10	30	.....	.....	149	120	0.163	.....	17.8	22.3
* Sample mostly lost in transit, and not included in average. † No records available, river levels as reported by collector.																		
STATION No. 14: FRASER RIVER AT																		
5	Sept. 9/50...	122	.....	.....	50	.....	..... (1.3)	7.7 (7.8)	10 (8)	15 (20)	11	11	122	73.6	0.100	.....	5.2	16.2
STATION No. 15: YELLOWHEAD																		
6	Sept. 9/50...	130	.....	.....	52	.....	.....	7.5 (8.2)	15 (5)	6 (5)	63	42	139	84.0	0.114	.....	25.2	16.4
STATION No. 16: PITT RIVER																		
7	Aug. 16/50...	92	Gauge height in feet		.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
			Max.	Min.														
			9.44	7.43														
			.....	.....	65	..... (9.9)	..... (1.5)	7.1 (7.1)	10 (10)	7 (5)	11	9.2	24.3	18.2	0.025	.....	5.4	2.6
STATION No. 17: ALOUETTE RIVER																		
8	Aug. 17/50...	69	.....	.....	483	65	..... (1.5)	7.0 (7.2)	0 (5)	0.5 .....	.....	.....	18.3	16.2	0.022	.....	5.2	3.2
Maximum flow for year, 13,000; minimum 21; average, 948 second-feet.																		
STATION No. 18: STAVE RIVER																		
9	Aug. 23/49...	38	.....	.....	62	.....	..... (2.0)	6.8 (6.8)	15 (5)	4 ( $<7$ )	2.4	1.2	15.2	22.8	0.031	.....	13.2	4.8
STATION No. 19: STAVE RIVER FROM																		
10	Aug. 17/50...	111	.....	.....	3,160	62	..... (2.0)	6.8 (6.7)	7 (5)	5 .....	.....	.....	13.4	.....	.....	.....	.....	2.2

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation Index		No.	
	Sodium (Na)	Potassium (K)	Total	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total			+	-		
BRIDGE NEAR TÊTE JAUNE CACHE—continued																					
5.8	1.8	0.9	.....	.....	16.5	0	.....	.....	0	77.3	0	.....	4.6	11.6	75.0	88.1	.....	.....	.....	1	
6.2	1.3	0.8	.....	.....	15.8	0	.....	.....	.....	81.0	0	.....	4.6	12.1	78.5	89.9	.....	.....	.....	2	
6.6	1.5	0.9	.....	0.17	17.9	0	0.4	0.05	.....	93.2	0	6.2	5.1	12.1	88.5	103	.....	.....	.....	3	
5.5	1.3	0.9	.....	0.15	16.5	0	0.5	0.04	.....	76.1	0	.....	3.8	15.9	78.3	88.1	3.4	.....	0.4	4	
BRIDGE NEAR MOUNT ROBSON																					
4.9	1.5	0.4	0.70	0.03	12.2	0	0.4	0.05	.....	63.2 (58.6)	0 (0)	.....	4.2	8.8	60.6	70.9	5.1	.....	0.8	5	
LAKE NEAR LUCERNE																					
7.9	0.9	0.3	0.41	0.02	16.5	0	0.7	0	.....	68.6 (65.9)	0 (0)	.....	2.7	17.3	73.5	79.1	2.6	.....	1.0	6	
FROM BRIDGE NEAR MOUTH																					
0.2	0.6	0.3	0.68	0.10	3.6	0	0.5	0.05	.....	9.8 (7.3)	0 (0)	.....	3.0	0	7.3	25.7	14.5	.....	3.0	7	
AT BRIDGE NEAR HANEY																					
0.1	0.6	0.1	0.10	.....	3.1	0	0	0	.....	8.5 (6.1)	0 (0)	3.4	3.7 (3.4)	1.4 (2.0)	8.4 (7.0)	15.0	13.3	.....	3.1	8	
AT STAVE FALLS																					
0	0.9	0.2	0.11	0.03	10.0	0	0.35	0.10	.....	5.4 (6.1)	0 (0)	4.6	3.0	7.6	12.0	21.9	13.8	.....	3.3	9	
HIGHWAY BRIDGE NEAR RUSKIN																					
0.1	0.5	0.2	.....	.....	2.5	0	.....	.....	.....	6.1	0	.....	2.7	0.9	5.9	11.2	15.0	.....	3.6	10	



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

## STATION No. 20: CANNELL

1	Aug. 23/49...	71	.....	.....	64	.....	.....	7.1 (6.4)	10 (10)	0.4	.....	.....	27.7	20.6	0.028	.....	8.6	3.8
---	---------------	----	-------	-------	----	-------	-------	--------------	------------	-----	-------	-------	------	------	-------	-------	-----	-----

## STATION No. 21: SUMAS

No.	Date of collection	Storage period (Days)	Gauge height in feet		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			Max.	Min.							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
2	Aug. 10/50...	123	7.45	.....	70	.....	.....	7.4 (6.5)	20 (7.6)	12 (30)	23	17	168	111	0.151	.....	13.8	16.0

## STATION No. 22: CHILLIWACK RIVER AT BRIDGE,

No.	Date of collection	Storage period (Days)	Gauge height in feet.		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			Max.	Min.							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
3	Aug. 25/49*	69	.....	.....	54	.....	.....	7.2 (1.5)	5 (7.9)	3 (5)	.....	.....	69.8	51.2	0.069	.....	6.2	12.0
4	Feb. 1/50*	27	6.10	6.66	32	.....	.....	7.7	5	0.4	.....	.....	96.6	63.6	0.087	.....	8.6	15.0
5	Feb. 25.....	12	7.40	6.66	42	.....	.....	7.9	5	15	52	48	101	59.4	0.081	.....	8.2	14.2
6	Mar. 26.....	22	7.10	7.62	42	.....	.....	7.7	5	6	13.0	12.6	89.7	58.6	0.080	.....	3.8	14.8
7	Apr. 25.....	14	7.60	7.34	42	.....	.....	7.8	10	7	18	15	81.3	55.0	0.075	.....	5.0	13.0
8	May 26.....	6	9.55	8.82	44	.....	.....	7.5	15	105	.....	.....	56.3	.....	.....	.....	.....	12.3
9	June 26.....	44	9.50	10.84	48	.....	.....	7.3	15	15	37	35	48.7	37.0	0.050	.....	10.0	7.5
10	Aug. 10.....	118	7.45	7.57	62	.....	.....	7.7 (3.0)	5 (7.85)	7 (5)	.....	.....	77.0	.....	.....	.....	.....	12.4
11	Aug. 15*.....	81	8.75	7.57	54	.....	.....	7.4	15	105	141	135	54.2	39.2	0.053	.....	9.4	8.1
12	Sept. 1.....	95	6.75	6.35	56	.....	.....	7.4	0	4	.....	.....	61.1	.....	.....	.....	.....	9.8
13	Sept. 13*.....	168	6.20	6.35	55	.....	.....	7.6	7	1	.....	.....	67.2	.....	.....	.....	.....	10.5
14	Sept. 25.....	70	6.65	6.35	52	.....	.....	7.6	3	5	.....	.....	67.9	.....	.....	.....	.....	10.6
15	Oct. 25.....	89	8.20	7.67	48	.....	.....	7.7	15	50	93	89	65.3	47.2	0.064	.....	6.0	10.4
16	Nov. 25.....	65	8.70	8.05	44	.....	.....	7.4	15	45	.....	.....	75.2	.....	.....	.....	.....	13.4
17	Dec. 12*.....	52	9.90	9.00	43	.....	.....	7.8	10	45	.....	.....	63.6	.....	.....	.....	.....	9.3
18	Dec. 26.....	43	10.45	9.00	43	.....	.....	7.4	15	35	.....	.....	65.2	.....	.....	.....	.....	10.3
19	Jan. 25/51...	19	8.05	7.13	38	.....	.....	7.4	10	25	49	46	83.5	57.4	0.078	.....	9.2	13.8
20	Average..... (12 samples)	49.8	8.1	7.8	46.8	.....	.....	7.6	9	27	.....	.....	72.7	52.4	0.071	.....	7.0	11.9

\* Not included in average.

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total <sup>b</sup>			+	-		
LAKE NEAR MISSION CITY																					
0.4	1.1	0.1	.....	0.05	6.6	0 (0)	0	0	.....	9.4 (11.0)	0 (0)	3.4	4.2	3.4	11.2	20.9	17.5	.....	2.9	1	
RIVER NEAR KILGARD																					
6.2	6.7	1.2	2.3	0.26	7.6	8.0	3.5	0.10	0.01	75.6 (79.3)	0 (0)	.....	21.0	3.4	65.4	108	17.8	.....	1.0	2	
VEDDER CROSSING—Drainage area, 450 square miles																					
0.9	1.6	0.5	.....	0.15	11.5	0	0.4	0	.....	25.4 (34.2)	0 (0)	4.4	7.2	12.8	33.6	46.8	.....	.....	.....	3	
2.4	1.8	0.3	.....	0.05	10.5	0	0.6	0	.....	45.4	0	5.6	7.3	10.1	47.3	.....	.....	.....	.....	4	
1.3	0.8	0.3	2.6	0.06	13.8	0	0.7	.....	.....	45.9	0	5.8	6.6	3.1	40.7	.....	.....	.....	.....	5	
1.3	1.4	0.6	1.1	0.03	13.5	0	0.4	0.05	.....	45.9	0	6.2	6.6	4.6	42.2	.....	.....	.....	.....	6	
0.8	1.7	0.6	1.0	0.04	11.0	0	0.5	0.05	.....	39.5	0	6.2	6.2	3.3	35.7	.....	.....	.....	.....	7	
0.8	0.9	0.5	.....	.....	12.8	0	.....	.....	.....	29.0	0	.....	5.0	10.2	34.0	.....	.....	.....	.....	8	
0.6	0.8	0.4	1.8	0.12	4.1	0	0.4	0	.....	19.5	0	.....	5.0	5.2	21.2	.....	.....	.....	.....	9	
1.0	1.4	0.4	.....	.....	7.8	0 (0)	.....	.....	.....	36.6 (36.6)	0 (0)	.....	6.2	5.0	35.0	.....	.....	.....	.....	10	
0.8	0.9	0.4	0.58	0.05	3.1	0	0.6	.....	.....	29.3	0	.....	4.3	0	23.5	.....	.....	.....	.....	11	
0.7	0.9	0.4	.....	.....	6.6	0	.....	.....	0	28.1	0	.....	6.1	4.4	27.4	.....	.....	.....	.....	12	
1.1	1.4	0.6	.....	.....	8.0	0	.....	.....	.....	34.2	0	.....	7.8	2.5	30.5	.....	.....	.....	.....	13	
0.9	1.1	0.4	.....	.....	4.9	0	.....	.....	.....	35.4	0	.....	5.7	1.1	30.1	.....	.....	.....	.....	14	
1.1	1.5	0.5	5.0	0.12	5.8	0	0.4	0.10	.....	34.4	0	.....	4.0	2.3	30.5	.....	.....	.....	.....	15	
1.1	2.2	1.4	.....	.....	4.5	0	.....	.....	0.01	48.8	0	.....	5.0	0	38.0	.....	.....	.....	.....	16	
1.1	0.9	0.6	.....	.....	6.8	0	.....	.....	.....	31.7	0	.....	5.0	1.5	27.5	.....	.....	.....	.....	17	
1.0	1.7	0.9	.....	.....	5.5	0	.....	.....	.....	35.6	0	.....	5.9	0.8	30.0	.....	.....	.....	.....	18	
1.4	0.9	0.2	2.5	0.10	4.9	0	0.5	.....	.....	42.0	0	5.4	5.0	5.8	40.2	.....	.....	.....	.....	19	
1.0	1.3	0.5	.....	0.08	7.9	0	0.8	0.06	0.01	36.7	0	.....	5.6	3.8	33.8	47.2	7.5	.....	1.2	20	



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>3</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

## STATION No. 23: CULTUS

1	Aug. 10/50....	118	.....	.....	73	.....	.....	8.0 (2.0)	5 (5)	3 .....	.....	.....	166	104	0.142	.....	7.2	29.8
---	----------------	-----	-------	-------	----	-------	-------	--------------	----------	------------	-------	-------	-----	-----	-------	-------	-----	------

## STATION No. 24: ELK

2	Aug. 25/49....	34	.....	.....	52	.....	.....	8.1 (1.2)	3 (5)	0.3 .....	.....	.....	238	154	0.209	.....	24.6	41.3
---	----------------	----	-------	-------	----	-------	-------	--------------	----------	--------------	-------	-------	-----	-----	-------	-------	------	------

## STATION No. 25: HARRISON RIVER

			Gauge height in feet															
			.....	.....														
3	Aug. 26/49*...	76	.....	.....	59	.....	.....	7.5 (1.5)	5 (5)	5 (7)	3.8	3.2	47.2	36.0	0.049	.....	7.8	7.6
4	Feb. 7/50....	25	4.53	4.78	37	.....	.....	7.5	5	2	.....	.....	46.8	38.4	0.052	.....	7.8	7.0
5	Mar. 6**....	10	5.85	4.13	40	.....	.....	7.3	0	4	5.0	3.2	48.8	34.0	0.046	.....	5.6	7.0
6	Apr. 3.....	14	3.50	4.57	42	.....	.....	7.6	0	0.5	.....	.....	49.1	33.0	0.045	.....	4.0	7.2
7	May 4.....	11	7.70	11.89	45	.....	.....	7.7	0	2	.....	.....	49.4	35.4	0.048	.....	7.6	7.0
8	June 5.....	31	15.16	18.94	51	.....	.....	7.4	3	0.4	.....	.....	49.4	.....	.....	.....	6.8	.....
9	July 3.....	37	19.48	16.64	51	.....	.....	7.3	5	3	.....	.....	50.8	38.0	0.052	.....	6.8	7.4
10	Aug. 1.....	79	13.65	11.03	58	.....	.....	7.4	6	9	.....	.....	45.1	.....	.....	.....	.....	7.2
11	Aug. 17*....	111	10.65	11.03	63	.....	.....	7.0 (1.3)	7 (5)	6 (5)	.....	.....	44.6	.....	.....	.....	6.7	.....
12	Sept. 5.....	40	9.35	8.01	60	.....	.....	7.2	2	5	.....	.....	43.8	.....	.....	.....	7.2	.....
13	Oct. 3.....	63	6.74	7.24	55	.....	.....	7.9	10	1	.....	.....	43.1	30.8	0.042	.....	5.0	6.7
14	Nov. 6.....	66	7.53	7.06	48	.....	.....	7.8	5	2	.....	.....	75.4	.....	.....	.....	10.5	.....
15	Dec. 3.....	61	6.97	7.62	45	.....	.....	7.2	5	3	.....	.....	45.1	.....	.....	.....	6.2	.....
16	Jan. 3.....	40	7.21	5.34	44	.....	.....	7.2	5	5	4.9	3.6	45.4	33.8	0.046	.....	9.0	6.2
17	Average..... (12 samples)	43.5	8.97	8.93	48	.....	.....	7.4	4	3	.....	.....	49.3	20.2	0.027	.....	6.5	7.2

\* Not included in average.

\*\* Heavy rains caused flood run-off from Chehalis River which enters above the sampling point.

## STATION No. 26: COQUIHALLA

18	Aug. 10/50....	132	.....	.....	57	.....	.....	7.4 (1.5)	3 (5)	1 (clear)	.....	.....	77.7	55.8	0.076	.....	12.0	13.4
----	----------------	-----	-------	-------	----	-------	-------	--------------	----------	--------------	-------	-------	------	------	-------	-------	------	------

TABLE II—Continued

Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-	

## LAKE AT CULTUS LAKE

2-3	3-2	0-5	.....	0-1	21-6	0 (0)	1-2	0	.....	73-2 (73-2)	1-2 (0)	8-0	6-2	21-0	83-9	102	7-8	.....	0-15	1
-----	-----	-----	-------	-----	------	----------	-----	---	-------	----------------	------------	-----	-----	------	------	-----	-----	-------	------	---

## RIVER NEAR CHILLIWACK

3-0	3-3	0-4	.....	0-04	39-5	1-6	1-2	0-05	.....	90-3 (95-2)	2-4 (0)	.....	10-9	37-3	115	148	5-8	0-1	.....	2
-----	-----	-----	-------	------	------	-----	-----	------	-------	----------------	------------	-------	------	------	-----	-----	-----	-----	-------	---

AT RAILWAY BRIDGE, HARRISON MILLS

0.5	1.3	0.6	0.06	0.02	7.9	0 (0)	0			19.9 (17.1)	0 (0)	6.4	6.4	4.8	21.1					3
0.5	1.4	0.7		0.06	6.3	0	0	0.05	0.02	21.5	0	4.6	5.8	8.1	19.6					
0.9	1.3	0.6	0.32	0.06	9.4	0	0	0.05		20.7	0	5.8	5.0	4.2	21.2					5
0.5	1.3	0.6		0.06	10.7	0	0	0.05		21.0	0	3.8	4.9	2.9	20.1					6
0.3	1.5	1.0		0.03	6.2	0	0.4	0.10	0	24.6	0	4.0	4.8	0	18.7					7
0.3	1.6	0.9			6.4	0				21.7	0		5.0	0.4	18.2					8
1.2	1.3	0.6		0.13	6.1	0.6	0.4	0.05		23.7	0	5.6	5.2	4.0	23.4					9
0.3	1.4	0.6			8.4	0			0	20.0	0		5.7	2.8	19.2					10
0.6	1.4	0.6			5.8	0				19.5 (17.1)	0 (0)		5.0	3.2	19.2					11
0.3	1.1	0.6			5.6	0			0	19.5	0		5.7	3.2	19.2					12
0.5	1.2	0.7		0.07	6.7	0.5	0.4	0		19.5	0		4.7	2.8	18.8					13
1.6	1.7	0.8			8.2	0	0		0.01	22.0	0		5.6	3.6	21.6					14
0.9	1.7	0.7			6.2	0				22.0	0		5.2	1.0	19.0					15
0.8	1.8	0.6	0.4	0.05	7.9	0	0	0.10		20.5	0		5.0	2.0	18.8					16
0.7	1.4	0.7		0.07	7.3	0	0.1	0.06		21.4	0		5.2	3.3	20.8	33.3	12.3		1.8	17

## RIVER NEAR HOPE

1.2	1.7	0.6	.....	0.10	8.2	0	0	0.05	.....	41.5 (36.6)	5 (0)	10.2	8.0	4.3	38.3	53.6	8.4	.....	1.3	18
-----	-----	-----	-------	------	-----	---	---	------	-------	----------------	----------	------	-----	-----	------	------	-----	-------	-----	----

TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>4</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C. (Ca)	Calcium
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 27: SCHKAM

1	Aug. 17/50....	69	.....	.....	67.5	.....	.....	7.2 (3.0)	3 (7.4)	0 (<5)	.....	.....	67.5	42.6	0.058	.....	5.4	8.6
---	----------------	----	-------	-------	------	-------	-------	--------------	------------	-----------	-------	-------	------	------	-------	-------	-----	-----

STATION No. 28: LYTTON

2	Aug. 18/50....	68	.....	.....	60	.....	.....	7.6 (3.0)	2 (7.8)	0.2 (<5)	.....	.....	132	87.6	.....	.....	5.4	22.3
---	----------------	----	-------	-------	----	-------	-------	--------------	------------	-------------	-------	-------	-----	------	-------	-------	-----	------

STATION No. 29: THOMPSON RIVER AT

			Gauge height in feet															
			.....	.....														
3	Aug. 27/49*....	75	.....	.....	64	.....	.....	7.5 (1.5)	5 (8.2)	4 (5)	8.6	7.2	86.3	56.8	0.077	.....	22.8	11.4
4	Mar. 1/50....	8	9,230	9,380	35	.....	.....	7.8	0	0.5	.....	.....	147	95.6	0.130	2.460	8.8	19.2
5	Mar. 28.....	8	8,510	9,380	40	.....	.....	8.1	0	0.3	.....	.....	130	78.0	0.106	1.779	10.0	16.6
6	Apr. 29.....	16	11,700	9,620	48	.....	.....	7.9	10	2	.....	.....	134	87.2	1.185	2.751	10.4	18.0
7	May 31.....	36	48,300	30,300	55	.....	.....	7.8	7	15	.....	.....	114	.....	.....	.....	.....	15.6
8	June—No sample taken.	.....	.....	84,300	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
9	July 4.....	36	91,600	68,400	62	.....	.....	7.5	7	15	28	27	85.3	59.0	0.080	1.456	9.6	12.1
10	Aug. 1.....	95	46,900	31,300	63	.....	.....	7.9	10	8	.....	.....	85.8	.....	.....	.....	.....	13.0
11	Aug. 9*.....	83	35,200	31,300	64	.....	.....	7.9 (1.5)	5 (7.9)	3 (5)	.....	.....	93.8	.....	.....	.....	.....	11.8
12	Sept. 6.....	83	20,900	18,300	65	.....	.....	7.6	2	5	.....	.....	88.3	.....	.....	.....	.....	12.5
13	Oct. 4.....	54	14,800	14,600	58	.....	.....	8.0	10	10	2.5	1.3	95.3	64.2	0.087	2.564	7.8	13.6
14	Oct. 31.....	94	13,200	14,600	48	.....	.....	7.4	8	1	.....	.....	104	74.4	0.101	2.650	9.0	15.0
15	Nov.—No sample taken.	.....	.....	12,900	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....
16	Dec. 1.....	69	12,400	11,600	37	.....	.....	7.8	5	0.5	.....	.....	102	.....	.....	.....	.....	14.2
17	Jan. 6/51....	38	10,400	9,330	35	.....	.....	7.7	10	0.8	.....	.....	83.4	.....	.....	.....	.....	14.6
18	Average..... (11 samples)	49	26,175	56,800	49	.....	.....	7.1	6	5	.....	.....	106.2	76.4	0.104	.....	9.3	14.9

\* Not included in average.

STATION No. 30: THOMPSON

19	Aug. 8/50....	84	35,800†	31,300†	62	.....	.....	7.8 (2.0)	8 (8.0)	6 (6)	.....	.....	86.5	.....	.....	.....	.....	11.5
----	---------------	----	---------	---------	----	-------	-------	--------------	------------	----------	-------	-------	------	-------	-------	-------	-------	------

† Records at Spences Bridge—see Station No. 29.



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.
	Sodium (Na)	Potassium (K)	Totale	Disolved								Gravi- metric	Colori- metric	Non- car- bonate	Totalb					
																		+	-	

## CREEK AT HOPE

1.7	1.2	0.8	.....	0.05	4.6	0 (0)	0.4	0	.....	37.3 (35.4)	0 (0)	5.4	7.0	0 (0)	28.5 (29.0)	42.7	8.1	.....	1.8	1
-----	-----	-----	-------	------	-----	----------	-----	---	-------	----------------	----------	-----	-----	----------	----------------	------	-----	-------	-----	---

## CREEK AT LYTTON

1.6	2.6	0.2	.....	0.03	5.3	0 (0)	0	0	.....	79.3 (78.1)	0 (0)	12	14	0 (1.0)	62.2 (65.0)	84.9	8.3	.....	0.7	2
-----	-----	-----	-------	------	-----	----------	---	---	-------	----------------	----------	----	----	------------	----------------	------	-----	-------	-----	---

## BRIDGE AT SPENCES BRIDGE—Drainage area, 21,500 square miles

1.8	1.4	0.9	0.34	0.02	6.8	0	0	0.05	.....	40.5	0	3.6	4.4	2.6	35.8	.....	.....	.....	.....	3
4.0	4.8	1.0	.....	.....	17.3	0	1.0	0.10	0.10	71.7	0	8.6	7.3	5.6	64.4	.....	.....	.....	.....	4
4.0	3.1	1.0	.....	0.02	20.9	0	0.4	0.05	.....	54.7	3.4	6.4	7.0	7.5	57.9	.....	.....	.....	.....	5
3.1	4.1	1.4	.....	0.06	21.3	0	0.5	0.15	.....	63.4	0	7.6	7.4	1.7	57.7	.....	.....	.....	.....	6
2.7	3.2	1.2	.....	.....	8.8	0	.....	.....	0.04	61.2	0	.....	9.2	0	50.0	.....	.....	.....	.....	7
.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	8
2.1	1.8	0.8	2.0	0.08	6.4	1.1	0	0.05	.....	47.3	0	.....	5.6	0	38.8	.....	.....	.....	.....	9
1.7	2.6	0.9	.....	.....	6.6	0	.....	.....	0.01	43.9	0	.....	9.3	3.4	39.4	.....	.....	.....	.....	10
1.7	1.8	0.8	.....	.....	7.0	0	.....	.....	.....	43.9	0	.....	5.0	0.4	36.4	.....	.....	.....	.....	11
2.0	1.9	0.9	.....	.....	8.6	0	.....	.....	0	45.4	0	.....	5.8	2.2	39.4	.....	.....	.....	.....	12
2.2	2.2	0.9	0.3	0.02	9.4	0	0.4	0.05	.....	48.8	0	.....	8.0	3.0	43.0	.....	.....	.....	.....	13
2.6	2.7	1.0	.....	0.20	11.0	0	0.7	0.10	.....	51.7	0	.....	7.0	5.6	48.0	.....	.....	.....	.....	14
.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	.....	15
2.5	2.2	0.8	.....	.....	8.3	0	.....	.....	0.05	53.7	0	.....	6.8	1.5	45.5	.....	.....	.....	.....	16
2.9	1.8	0.8	.....	.....	12.2	0	.....	.....	.....	54.2	0	.....	4.8	4.0	48.4	.....	.....	.....	.....	17
2.7	2.8	1.1	.....	0.07	11.9	.....	0.6	0.08	0.05	54.2	.....	.....	7.1	3.9	48.3	68.0	10.9	.....	1.4	18

## RIVER AT ASHCROFT

1.6	1.8	0.8	.....	.....	5.8	0	.....	.....	.....	41.7	0	.....	4.2	1.1 (5.0)	35.3 (37.0)	46.2	9.7	.....	0.9	16
-----	-----	-----	-------	-------	-----	---	-------	-------	-------	------	---	-------	-----	--------------	----------------	------	-----	-------	-----	----

TABLE II—Continued

Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 31: THOMPSON RIVER AT

			Gauge height in feet															
1	Aug. 27/49....	13	.....	.....	64	.....	.....	7.4 (1.5)	5 (8.1)	6 (8)	.....	.....	80.8	.....	.....	.....	.....	11.0
2	Aug. 8/50....	120	6.15†	5.74†	61	.....	.....	7.6 (2.0)	6 (7.8)	3 (7)	.....	.....	80.0	.....	.....	.....	.....	11.6

† Records at Kamloops, B.C.—see Station No. 33.

STATION No. 32: THOMPSON\* RIVER

3	Feb. 1-10/50.	32	2.74		33	.....	.....	7.5	5	1	.....	.....	19.1	122	0.166	.....	7.2	12.9
4	Feb. 11-20....	29	2.55	2.60	34	.....	.....	7.7	0	0.5	.....	.....	99.6	64.0	0.087	.....	8.2	13.2
5	Feb. 21-28....	20	2.50		33	.....	.....	7.5	0	0.5	.....	.....	101	65.2	0.089	.....	9.4	13.0
6	Mar. 1-10....	31	2.00		34	.....	.....	7.6	0	0.3	.....	.....	109	66.2	0.090	.....	11.6	13.5
7	Mar. 11-20....	33	0.54	0.86	34	.....	.....	7.4	0	0.3	.....	.....	103	66.2	0.090	.....	6.0	14.2
8	Mar. 21-31....	32	0.14		40	.....	.....	7.6	2	2	.....	.....	101	82.6	0.112	.....	17.4	14.6
9	Apr. 1-10....	22	0.12		42	.....	.....	7.6	3	3	.....	.....	98.6	73.0	0.099	.....	11.4	14.6
10	Apr. 11-20....	30	0.41	0.46	44	.....	.....	7.9	0	1	.....	.....	107	69.2	0.094	.....	10.6	13.8
11	Apr. 21-30....	20	0.86		44	.....	.....	7.6	8	0.7	.....	.....	101	70.4	0.096	.....	6.6	13.0
12	May 1-10....	32	1.86		48	.....	.....	7.8	5	0.3	.....	.....	95.5	66.8	0.091	.....	8.2	13.4
13	May 11-20....	36	5.27	4.64	52	.....	.....	7.5	5	0.4	.....	.....	93.6	65.4	0.089	.....	6.0	14.4
14	May 21-31....	25	6.59		48	.....	.....	7.4	0	0.4	.....	.....	90.1	66.0	0.090	.....	7.2	12.1
15	June 1-10....	37	8.32		50	.....	.....	7.5	2	4	7.0	3.4	90.1	113	0.154	.....	44.6	12.8
16	June 11-20....	19	13.42	12.51	54	.....	.....	7.4	8	0.3	.....	.....	92.7	66.2	0.090	.....	14.0	12.8
17	June 21-30....	6	15.78		57	.....	.....	7.4	3	0.5	.....	.....	82.0	64.6	0.088	.....	11.6	13.2
18	July 1-10....	71	13.44		59	.....	.....	7.4	5	0.4	.....	.....	83.8	62.8	0.085	.....	10.6	13.6
19	July 11-20....	71	10.66	10.52	58	.....	.....	7.6	5	3	.....	.....	82.0	69.8	0.095	.....	11.8	12.2
20	July 21-31....	101	8.70		60	.....	.....	7.4	5	1	.....	.....	93.1	61.4	0.084	.....	9.0	11.0

\* Due to sample location being on left side of bridge, water is probably usually from the South Thompson River.

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		

## HIGHWAY BRIDGE NEAR SAVONA—Drainage area at Walhachin, 15,600 square miles

1.4	1.6	0.9	.....	.....	4.9	0	.....	.....	.....	39.0 (41.5)	0 (0)	.....	5.2	1.3	33.3	44.2	9.2	.....	1.5	1
1.4	1.2	0.8	.....	.....	6.8	0	.....	.....	.....	38.1 (34.2)	0 (0)	.....	5.4	3.5	34.7	46.0	6.8	.....	1.2	2

## AT BRIDGE BELOW KAMLOOPS—Drainage area, 14,400 square miles

2.6	18.8	1.2	.....	0.27	11.4	25.8	0.4	0.10	0	51.0	0	12.4	9.9	1.1	42.9	.....	.....	.....	.....	3
2.3	2.4	1.0	.....	0.02	13.0	0	0.6	0.10	.....	46.6	0	7.4	6.2	4.3	42.5	.....	.....	.....	.....	4
2.5	2.0	0.9	.....	0.04	12.8	0	0.4	0.10	.....	45.9	0	7.2	6.3	5.1	42.7	.....	.....	.....	.....	5
2.5	2.4	0.9	.....	0.03	16.0	0	0.4	0.10	.....	46.6	0	7.6	7.3	5.6	44.0	.....	.....	.....	.....	6
2.5	2.3	0.9	.....	0.03	17.3	0	0.7	0.10	0	47.3	0	5.6	6.8	6.9	45.7	.....	.....	.....	.....	7
2.7	2.4	1.0	.....	0.07	14.0	0	0.6	0.15	.....	48.8	0	8.2	7.6	7.5	47.5	.....	.....	.....	.....	8
2.0	2.4	1.0	.....	0.17	18.0	0	0.8	0.10	0.01	47.6	0	9.4	7.0	9.7	48.7	.....	.....	.....	.....	9
2.3	3.2	0.9	.....	0.05	12.8	0	0.8	0.10	.....	50.0	0	6.4	6.8	2.9	43.9	.....	.....	.....	.....	10
2.2	2.4	1.1	.....	0.06	7.4	0	0.7	0.10	.....	46.4	0	6.4	6.7	3.5	41.5	.....	.....	.....	.....	11
1.9	2.1	0.9	.....	0.12	9.5	0	1.3	0.10	.....	45.9	0	7.8	7.0	3.6	41.2	.....	.....	.....	.....	12
2.1	1.8	1.0	.....	0.22	7.1	0	0.9	0.10	.....	47.8	0	11.0	7.3	5.4	44.6	.....	.....	.....	.....	13
1.9	1.6	0.9	.....	0.17	8.1	0	0.8	0.10	.....	44.9	0	11.4	5.6	1.2	38.0	.....	.....	.....	.....	14
1.8	2.0	1.0	0.8	0.12	8.9	0	0.5	0.05	0	42.2	0	.....	6.4	4.7	39.3	.....	.....	.....	.....	15
1.9	1.8	0.9	.....	0.13	8.4	0	1.2	0.08	.....	43.9	0	.....	9.4	3.7	39.7	.....	.....	.....	.....	16
1.7	2.0	0.9	.....	0.25	10.4	0	1.3	0.08	.....	41.5	0	.....	8.6	5.9	39.9	.....	.....	.....	.....	17
1.5	1.8	0.9	.....	0.16	8.6	0	0.5	0.10	.....	42.7	0	.....	8.8	5.1	40.1	.....	.....	.....	.....	18
1.7	1.8	0.8	.....	0.19	6.4	0	0.9	0.10	.....	42.0	0	.....	10.0	3.0	37.4	.....	.....	.....	.....	19
1.5	1.8	0.9	.....	0.17	4.9	0	0.8	.....	.....	43.9	0	.....	7.2	0	33.6	.....	.....	.....	.....	20



TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>4</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 32—THOMPSON RIVER AT

		Gauge height in feet																		
1	Aug. 1-10....	90	6.80	5.74	63			7.5	3	3				79.8	60.8	0.083		11.8	10.7	
2	Aug. 8/50*...	120	6.15		61			7.6	6	3				80.0					11.6	
3	Aug. 8/50**...	84	6.15		61		(2.0)	7.6	(7)	(5)				88.3					12.5	
4	Aug. 11-20....	92	5.47		64		(1.1)	7.7	(8)	(15)				78.6	56.2	0.076		8.8	11.4	
5	Aug. 21-31....	93	5.03	3.34	63			7.6	10	3				79.0	60.0	0.082		7.4	11.7	
6	Sept. 1-10....	97	4.21		63			7.2	4	5	7.4	3.2		80.8	55.0	0.075		8.0	11.0	
7	Sept. 11-20....	81	3.05		64			7.9	10	0.8				77.3	56.6	0.077		9.6	11.4	
8	Sept. 21-30....	63	2.75		61			7.6	0	3				80.1	59.4	0.081		8.8	11.7	
9	Oct. 1-10....	67	2.11	2.24	54			7.2	1	5	9.8	5.6		83.8	57.4	0.078		8.8	12.0	
10	Oct. 11-20....	110	2.74		52			7.5	7	0.4				82.8	63.2	0.086		9.0	12.1	
11	Oct. 21-31....	100	1.90		48			7.3	5	0.6				86.2	58.4	0.079		10.8	11.8	
12	Nov. 1-10....	109	2.04		44			7.4	5	0.7				84.3	60.8	0.083		12.8	13.0	
13	Nov. 11-20....	89	1.72	1.68	39			7.6	5	0.5				82.9	59.6	0.081		6.8	12.4	
14	Nov. 21-30....	73	1.29		38			7.3	5	3				86.6	63.0	0.086		11.0	12.5	
15	Dec. 1-10....	53	1.35		36			7.2	7	3				92.3	62.0	0.084		13.8	12.7	
16	Dec. 11-20....	59	1.11		37			7.0	3	1				90.8	60.8	0.083		13.2	12.5	
17	Dec. 21-31....	59	0.99	1.15	38			7.8	6	0.6				69.4	63.6	0.087		9.0	13.9	
18	Jan. 1-10/51.	52	0.99		36			7.3	5	0.5				95.8	66.4	0.090		11.4	13.1	
19	Jan. 11-20....	42	0.85		0.96	33.5			7.2	5	0.5				99.2	69.4	0.094		14.2	13.7
20	Jan. 21-31....	43	1.05			33			7.2	10	0.3				97.6	64.0	0.087		12.4	13.3
21	Average, 12 mos. (36 samples)	56	3.95	3.90		46.7			7.5	4	5				92.8	67.0	0.091		11.1	12.8

\* Left bank—South Thompson River  
\*\* Right bank—North Thompson River

STATION No. 33: NORTH THOMPSON

22	Aug. 27/49....	75			62			7.6	15	7	20	17		82.3	47.0	0.064		22.0	11.6
							(2.0)	(8.0)		(10)									

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		

## BRIDGE BELOW KAMLOOPS—Continued

1.7	1.9	0.8	.....	0.16	5.8	0	0.4	.....	0.01	39.5	0	.....	6.7	1.3	33.7	.....	.....	.....	.....	.....	1
1.4	1.2	0.8	.....	.....	6.8	0	.....	.....	.....	38.1 (34.2)	0 (0)	.....	5.4	3.5	34.7	.....	.....	.....	.....	.....	2
1.5	1.3	0.8	.....	.....	6.6	0	.....	.....	.....	43.9	0	.....	3.4	1.4 (6.0)	37.4 (38.0)	.....	.....	.....	.....	.....	3
1.6	1.8	0.6	0.46	0.03	6.9	0	0.5	0.10	.....	39.8	0	.....	8.3	2.4	35.0	.....	.....	.....	.....	.....	4
1.4	2.0	0.9	.....	0.08	6.6	0	0.5	0.05	.....	41.0	0	.....	10.6	1.4	35.0	.....	.....	.....	.....	.....	5
1.8	0.7	1.2	0.4	0.02	6.6	0	1.3	0	.....	38.6	0	.....	7.6	3.2	34.8	.....	.....	.....	.....	.....	6
1.6	2.0	0.9	.....	0.07	9.5	0.5	0.9	0.10	.....	39.3	0	5.2	7.5	2.8	35.0	.....	.....	.....	.....	.....	7
1.7	1.9	0.9	.....	0.12	7.4	0	0.5	0.10	.....	41.0	0	.....	9.0	2.6	36.2	.....	.....	.....	.....	.....	8
2.0	1.7	1.2	0.6	0.02	9.1	0	1.1	0.10	.....	40.7	0	.....	6.3	4.8	38.2	.....	.....	.....	.....	.....	9
1.9	2.2	0.7	.....	0.12	7.9	0	0.7	0.10	.....	39.8	0	.....	7.2	5.4	38.0	.....	.....	.....	.....	.....	10
1.9	2.1	0.7	.....	0.16	6.6	0	1.3	0.10	.....	43.4	0	.....	6.4	1.7	37.3	.....	.....	.....	.....	.....	11
1.8	1.9	1.0	.....	0.01	6.6	0	0	.....	0.05	43.4	0	.....	3.2	4.3	39.9	.....	.....	.....	.....	.....	12
2.0	1.3	0.8	.....	0.14	8.9	0	0.6	0.10	.....	43.7	0	.....	8.0	3.4	39.2	.....	.....	.....	.....	.....	13
2.1	2.5	1.1	.....	0.10	8.2	0	1.8	0.05	.....	43.9	0	.....	8.0	3.8	39.8	.....	.....	.....	.....	.....	14
2.1	2.2	0.9	.....	0.12	6.4	0	1.3	0.10	.....	46.4	0	.....	5.8	2.3	40.3	.....	.....	.....	.....	.....	15
2.1	2.0	0.9	.....	0.08	9.5	0	0.9	0.10	.....	43.9	0	.....	6.2	3.8	39.8	.....	.....	.....	.....	.....	16
2.5	2.0	1.0	.....	0.02	13.2	0	0.9	.....	.....	46.4	0	.....	3.6	7.0	45.0	.....	.....	.....	.....	.....	17
2.1	1.9	0.9	.....	0.07	7.0	0	2.2	0.10	.....	46.1	0	6.6	4.3	3.5	41.3	.....	.....	.....	.....	.....	18
2.1	1.9	0.8	.....	0.06	8.6	0	1.3	0.10	.....	46.1	0	6.2	5.0	5.0	42.8	.....	.....	.....	.....	.....	19
2.3	2.1	0.8	.....	0.08	10.2	0	1.3	0	.....	45.9	0	8.2	6.4	5.1	42.7	.....	.....	.....	.....	.....	20
2.0	2.5	0.7	.....	0.10	9.4	0	0.86	0.08	.....	44.2	0	.....	7.1	4.0	40.2	50.1	11.7	.....	.....	1.2	21

## RIVER AT RAYLEIGH

1.8	1.3	1.0	1.2	0.01	6.3	0 (0)	0.9	0.05	.....	38.2 (37.8)	0 (0)	2.6	2.4	5.0	36.3	44.1	7.0	.....	.....	1.3	22
-----	-----	-----	-----	------	-----	----------	-----	------	-------	----------------	----------	-----	-----	-----	------	------	-----	-------	-------	-----	----

TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>3</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 34: NORTH THOMPSON RIVER AT

		Gauge height in feet																
1	Feb. 20/50....	12	2,760	2,620	36	.....	7.8	5	0.5	.....	.....	.....	107	71.8	0.098	0.534	10.2	15.6
2	Mar. 20.....	7	2,760	2,810	37	.....	7.7	5	0.4	.....	.....	.....	116	73.0	0.099	0.544	6.4	16.8
3	Apr. 20.....	14	3,780	3,600	46	.....	7.9	5	3	.....	.....	.....	101	71.0	0.097	0.724	13.6	14.2
4	May 20.....	10	24,800	19,100	43	.....	7.8	20	5	.....	.....	.....	77.4	.....	.....	.....	.....	12.0
5	June 20.....	22	74,100	52,700	53	.....	7.2	20	15	.....	.....	.....	61.7	45.4	0.062	9.08	7.4	9.0
6	July 21.....	84	33,600	35,800	51	.....	7.5	5	25	45	42	.....	68.9	46.2	0.063	4.19	10.2	11.5
7	Aug. 7*.....	85	18,300	17,800	62	.....	7.6 (1.5)	8 (7.6)	10 (5)	.....	.....	.....	82.7	.....	.....	.....	.....	11.4
8	Aug. 19.....	77	15,800	17,800	58	.....	7.4	5	9	.....	.....	.....	77.2	.....	.....	.....	.....	12.3
9	Sept. 21.....	64	8,000	13,500	55	.....	7.3	2	10	.....	.....	.....	86.1	.....	.....	.....	.....	12.4
10	Oct. 21.....	92	6,080	7,200	48	.....	7.6	3	5	11	98	.....	90.4	62.2	0.084	1.01	10.8	13.5
11	Nov. 22.....	54	3,700	5,620	34	.....	7.2	5	3	.....	.....	.....	119	.....	.....	.....	.....	16.3
12	Dec. 21.....	43	3,780	4,020	36	.....	7.4	10	15	.....	.....	.....	102	.....	.....	.....	.....	18.8
13	Jan. 22/51....	22	2,810	3,070	37	.....	7.4	7	0.5	.....	.....	.....	116	75.4	0.103	0.573	10.4	17.6
14	Average..... (12 samples)	42	15,122	13,987	44.5	.....	7.6	8	9	.....	.....	.....	93.5	63.6	0.087	.....	9.8	14.2

\* Field sample, not included in average.

STATION No. 35: NORTH THOMPSON RIVER

15	Sept. 5/50....	125	Fast	.....	54 (12.0)	.....	7.6 (1.5)	20 (7.8)	85 (4.5)	53	52	.....	53.3	40.4	0.055	.....	4.0	7.0
----	----------------	-----	------	-------	--------------	-------	--------------	-------------	-------------	----	----	-------	------	------	-------	-------	-----	-----

STATION No. 36: CLEARWATER RIVER AT

16	Sept. 5/50....	106	Fast	.....	59	.....	7.6 (2.0)	5 (8.0)	3 (10)	3 (< 5)	.....	.....	89.8	56.4	0.077	.....	7.4	16.2
----	----------------	-----	------	-------	----	-------	--------------	------------	-----------	------------	-------	-------	------	------	-------	-------	-----	------

STATION No. 37: RAFT RIVER AT

17	Sept. 5/50....	106	.....	.....	57	.....	7.6 (2.0)	10 (8.0)	3 (15)	.....	.....	.....	78.6	56.4	0.077	.....	7.4	12.2
----	----------------	-----	-------	-------	----	-------	--------------	-------------	-----------	-------	-------	-------	------	------	-------	-------	-----	------



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total			+	-		

## HIGHWAY BRIDGE, BARRIERE—Drainage area, 7,040 square miles

2.6	2.0	1.1	.....	0.03	10.0	0	0.7	0.05	0.06	53.7	0	6.6	7.0	5.6	49.6	.....	.....	.....	.....	.....	1
3.3	2.4	1.1	.....	0.15	15.3	0	0.5	0.05	.....	58.8	0	7.2	6.7	7.3	55.5	.....	.....	.....	.....	.....	2
2.3	2.2	1.4	.....	0.25	12.2	0	0	0.05	.....	51.2	0	8.8	6.0	2.9	44.9	.....	.....	.....	.....	.....	3
1.8	0.9	0.7	.....	.....	10.5	0	.....	.....	0	41.5	0	.....	5.2	3.3	37.3	.....	.....	.....	.....	.....	4
1.0	1.3	0.8	2.0	0.07	5.8	0	0.9	0.01	.....	31.7	0	.....	3.7	0.6	26.6	.....	.....	.....	.....	.....	5
1.0	0.9	1.0	2.3	0.12	6.1	0	0.8	.....	.....	36.6	0	.....	4.2	2.8	32.8	.....	.....	.....	.....	.....	6
1.2	1.3	0.8	.....	.....	6.2	0	.....	.....	.....	39.3	0	.....	3.1	1.1	33.3	.....	.....	.....	.....	.....	7
1.2	0.9	0.8	.....	.....	8.6	0	.....	.....	0.01	36.8	0	.....	5.2	5.4	35.6	.....	.....	.....	.....	.....	8
1.7	0.9	0.9	.....	.....	8.6	0	.....	.....	.....	42.0	0	.....	4.1	3.5	37.9	.....	.....	.....	.....	.....	9
1.9	1.3	1.0	0.6	0.05	8.2	0	0.5	0.05	.....	47.1	0	.....	5.6	2.9	41.5	.....	.....	.....	.....	.....	10
2.1	1.8	1.2	.....	.....	9.1	0	.....	.....	0.02	53.7	0	.....	3.2	5.5	49.5	.....	.....	.....	.....	.....	11
2.5	1.8	1.1	.....	.....	15.6	0	.....	.....	.....	52.7	0	.....	4.0	13.8	57.0	.....	.....	.....	.....	.....	12
2.7	1.8	1.1	.....	0.09	9.7	0	0.8	0.10	.....	61.0	0	6.6	6.4	5.0	55.0	.....	.....	.....	.....	.....	13
2.0	1.6	1.0	.....	0.11	10.0	0	0.6	0.05	.....	47.1	0	.....	5.1	5.1	43.7	57.9	4.4	.....	.....	1.0	14

## AT HIGHWAY BRIDGE, CLEARWATER

1.5	1.8	1.6	5.1	0.44	7.4	0	0.5	0.05	.....	23.9 (22.0)	0 (0)	.....	4.8	4.0	23.6	36.7	13.2	.....	.....	1.7	15
-----	-----	-----	-----	------	-----	---	-----	------	-------	----------------	----------	-------	-----	-----	------	------	------	-------	-------	-----	----

## HIGHWAY BRIDGE, CLEARWATER

1.9	0.9	0.7	.....	0.08	5.8	0	1.2	0	.....	48.8 (46.6)	0 (0)	4.0	1.9	8.2	48.2	52.1	4.2	.....	.....	1.0	16
-----	-----	-----	-------	------	-----	---	-----	---	-------	----------------	----------	-----	-----	-----	------	------	-----	-------	-------	-----	----

## HIGHWAY BRIDGE NEAR CLEARWATER

1.9	0.9	0.7	.....	0.08	5.0	0	0.7	0.05	.....	48.8	0	6.4	9.6	0	38.2	55.1	4.6	.....	.....	1.2	17
-----	-----	-----	-------	------	-----	---	-----	------	-------	------	---	-----	-----	---	------	------	-----	-------	-------	-----	----

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>3</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

## STATION No. 38: SOUTH THOMPSON RIVER

		Gauge height in feet																
No.	Date	Storage period (Days)	On sampling date	Monthly mean	Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Dried at 105°C.	Ignited at 550°C.	Specific conductance K x 10 <sup>3</sup> at 25°C.	P.P.M.	Tons per acre-foot	Thousand tons per day	Loss on ignition at 550°C.	Calcium (Ca)
1	Feb. 20/50....	12	3,350	3,560	33	.....	.....	7.7	5	0.4	.....	.....	82.4	56.0	0.077	0.510	8.2	13.0
2	Mar. 20.....	7	3,000	3,040	39	.....	.....	7.7	5	0.3	.....	.....	83.7	55.6	0.076	0.450	6.6	12.5
3	Apr. 20.....	7	3,620	3,490	43	.....	.....	7.8	5	0.5	.....	.....	78.8	54.0	0.0735	0.528	9.0	13.4
4	May 20.....	10	11,700	9,540	49	.....	.....	7.8	5	0.4	.....	.....	85.5	.....	.....	.....	.....	12.8
5	June 20.....	16	34,000	28,200	49	.....	.....	7.5	5	0.5	.....	.....	79.2	.....	.....	.....	.....	11.0
6	June 26†.....	28	39,500	28,200	58	.....	.....	7.5	8	0.9	.....	.....	79.9	59.4	0.081	6.24	12.0	13.2
7	July 20.....	53	25,800	29,300	63	.....	.....	7.5	8	0.5	.....	.....	80.8	54.6	0.074	3.80	6.4	11.8
8	Aug. 7*.....	85	15,000	12,500	63	.....	.....	7.7 (1.5)	8 (7.8)	0.3 (5)	.....	.....	81.3	.....	.....	.....	.....	10.8
9	Aug. 31.....	65	9,140	12,500	65	.....	.....	7.6	5	0.2	.....	.....	74.0	.....	.....	.....	.....	11.2
10	Sept. 20.....	161	8,560	9,900	67	.....	.....	7.9	2	0.4	.....	.....	70.6	.....	.....	.....	.....	10.3
11	Oct. 20.....	46	4,990	5,060	47	.....	.....	7.6	2	4	.....	.....	71.3	50.6	0.069	0.682	8.8	11.2
12	Nov. 20.....	70	4,510	4,660	40	.....	.....	7.5	5	0.5	.....	.....	73.2	.....	.....	.....	.....	11.2
13	Dec. 20.....	44	4,050	4,110	38	.....	.....	7.5	6	15	.....	.....	79.8	.....	.....	.....	.....	12.8
14	Jan. 20/51....	24	3,940	4,030	34	.....	.....	7.2	5	0.4	.....	.....	79.5	54.0	0.735	0.574	8.8	12.5
15	Average..... (12 samples)	43	4,560	9,782	47	.....	.....	7.6	5	0.6	.....	.....	78.2	54.1	0.074	.....	7.9	11.9

† Flood sample not included in average.

\* Field sample not included in average.

## STATION No. 39: ADAMS RIVER AT

16	Aug. 5/50....	122	3,960	2,350	61	.....	.....	7.5 (0)	5 (8.0)	0.7 (5)	.....	.....	61.2	46.0	0.064	0.491	8.2	9.2
----	---------------	-----	-------	-------	----	-------	-------	------------	------------	------------	-------	-------	------	------	-------	-------	-----	-----

## STATION No. 40: SHUSWAP

17	Aug. 5/50....	67	.....	.....	61.5	.....	.....	8.1 (0)	7 (8.5)	6 (10)	.....	.....	123	82.8	0.113	.....	11.0	24.6
----	---------------	----	-------	-------	------	-------	-------	------------	------------	-----------	-------	-------	-----	------	-------	-------	------	------

## STATION No. 41: SHUSWAP

18	Aug. 4/50....	52	.....	.....	67	.....	.....	7.7 (1.5)	40 (8.0)	1 (<5)	.....	.....	86.2	63.4	0.086	.....	11.2	14.2
----	---------------	----	-------	-------	----	-------	-------	--------------	-------------	-----------	-------	-------	------	------	-------	-------	------	------

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.
	Sodium (Na)	Potassium (K)	Total	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total			+	-	

## AT HIGHWAY BRIDGE, CHASE—Drainage area, 6,060 square miles

1-7	1-4	0-9	.....	0-03	8-7	0	0-4	0	0-02	41-5	0	5-8	6-8	5-4	39-4	.....	.....	.....	.....	1
2-0	1-3	0-8	.....	0-03	7-4	0	0-4	0-10	.....	45-9	0	6-4	6-0	1-8	39-4	.....	.....	.....	.....	2
2-1	1-2	0-8	.....	0-05	16-6	0	0	0-10	.....	39-0	0	7-4	6-2	10-0	42-0	.....	.....	.....	.....	3
1-8	1-1	0-7	.....	.....	11-5	0	.....	.....	0-02	43-7	0	.....	6-1	3-5	39-3	.....	.....	.....	.....	4
0-8	1-6	0-9	.....	.....	6-6	0	.....	.....	.....	41-0	0	.....	6-4	0	30-7	.....	.....	.....	.....	5
1-6	1-6	0-9	.....	0-11	12-0	0	0-98	0-05	.....	39-3	0	.....	7-8	7-3	39-5	.....	.....	.....	.....	6
1-8	1-6	0-9	.....	0-05	6-4	0-2	1-0	0-10	.....	39-0	0	8-2	5-8	4-8	36-8	.....	.....	.....	.....	7
1-2	1-4	0-8	.....	.....	3-5	0	.....	.....	.....	42-9	0	.....	5-8	0	31-8 (30-0)	.....	.....	.....	.....	8
1-2	1-3	0-8	.....	.....	6-2	0	.....	.....	0-02	36-6	0	.....	7-1	2-8	32-8	.....	.....	.....	.....	9
1-6	1-7	0-9	.....	.....	5-3	0	.....	.....	.....	36-4	0	.....	6-7	2-7	32-5	.....	.....	.....	.....	10
1-3	1-2	0-7	.....	0-05	7-1	0	0-4	0	.....	37-3	0	4-2	5-7	2-6	33-2	.....	.....	.....	.....	11
2-0	1-7	1-1	.....	.....	6-2	0	.....	.....	0	41-5	0	.....	5-7	2-0	36-0	.....	.....	.....	.....	12
1-9	1-7	1-1	.....	.....	12-1	0	.....	.....	.....	42-5	0	.....	5-4	5-2	40-0	.....	.....	.....	.....	13
1-8	0-8	0-6	.....	0-04	5-9	0	< .4	0-10	.....	42-0	0	5-2	6-2	4-2	38-6	.....	.....	.....	.....	14
1-7	1-4	0-9	.....	0-05	8-3	0	0-4	0-07	.....	40-5	0	.....	6-2	3-5	36-7	51-0	7-6	.....	1-2	15

## HIGHWAY BRIDGE NEAR SQUILAX—Drainage area, 1,600 square miles

1-1	1-4	0-6	.....	0-06	6-6	0	0-5	0	.....	30-5	0	4-8	6-4	2-5	27-5	40-8	9-7	.....	1-5	16
-----	-----	-----	-------	------	-----	---	-----	---	-------	------	---	-----	-----	-----	------	------	-----	-------	-----	----

## LAKE AT SALMON ARM

3-0	2-7	1-1	0-42	0-07	9-7	0	0-3	0-10	.....	67-1 (61-1)	0 (4-8)	.....	9-3	18-7	73-7	84-0	9-0	.....	0-15	17
-----	-----	-----	------	------	-----	---	-----	------	-------	----------------	------------	-------	-----	------	------	------	-----	-------	------	----

## RIVER NEAR ENDERBY—Drainage area near Lumby, B.C.—650 square miles

1-4	0-8	0-7	.....	0-09	6-6	0	0-4	0	.....	46-4 (45-1)	0 (0)	.....	7-2	3-2 (5-0)	41-2 (42-0)	54-1	4-0	.....	0-9	18
-----	-----	-----	-------	------	-----	---	-----	---	-------	----------------	----------	-------	-----	--------------	----------------	------	-----	-------	-----	----



TABLE II—Continued

Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
STATION No. 42: MARA																		
1	Aug. 28/49....	12	.....	.....	70	.....	..... (2.0)	7.7 (8.2)	5 (5)	6	.....	.....	96.0	.....	.....	.....	16.0	
STATION No. 43: MARA																		
2	Aug. 4/50....	27	.....	.....	70	.....	..... (2.0)	7.7 (8.0)	8 (5)	1 ( $<5$ )	.....	.....	89.1	59.0	0.080	.....	9.4	16.4
STATION No. 44: SALMON RIVER FROM																		
3	Aug. 5/50....	103	.....	.....	62	.....	..... (0)	8.1 (8.0)	9 (6)	4 .....	7.0	5.7	359	23.9	0.325	.....	19.4	43.9
STATION No. 45: EAST CANOE CREEK NEAR																		
4	Aug. 5/50....	67	.....	.....	59	.....	..... (3.0)	8.0 (7.9)	5 (5)	0.7 .....	.....	.....	362*	213*	0.290	.....	9.8	67.7*
* Calcium corrected for loss on storage; conductivity and residue on evaporation low.																		
STATION No. 46: EAGLE																		
5	Aug. 4/50....	104	.....	.....	49	.....	.....	7.4 (7.6)	6 (5)	6 (5)	.....	.....	49.4	.....	.....	.....	.....	6.7
STATION No. 47: EAGLE																		
6	Aug. 29/49....	32	.....	.....	59	.....	.....	7.2	10	5	10.0	7.2	49.2	53.6	0.073	.....	21.8	9.2
STATION No. 48: NORTH FORK EAGLE																		
7	Aug. 4/50....	104	.....	.....	49	.....	..... (1.0)	7.5 (7.6)	5 (5)	6 (5)	8.2	6.6	33.6	26.0	0.035	.....	5.6	5.0
STATION No. 49: NICOLA RIVER AT																		
8	Aug. 8/50....	23	.....	.....	67	.....	..... (5.0)	8.5 (8.6)	10 (8.6)	3 .....	.....	.....	249	147	0.200	.....	17.0	30.8

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.	
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		
LAKE NEAR MARA																					
1.5	1.4	1.0	.....	.....	9.8	0	.....	.....	.....	48.6 (51.3)	0 (0)	.....	6.8	6.3	46.1	60.4	6.0	.....	0.9	1	
LAKE NEAR SICAMOUS																					
1.7	1.4	0.9	.....	0.05	11.0	0	0	0.05	.....	48.3 (46.5)	0 (0)	.....	3.3	8.3	47.9	23.7	5.8	.....	0.8	2	
HIGHWAY BRIDGE AT SALMON ARM																					
12.9	13.7	2.9	0.50	0.03	31.8	0	0.4	0.20	.....	194 (187)	4.8 (4.8)	.....	27	0 (13.0)	163 (174)	233	15.2	0.5	.....	3	
SALMON ARM—Drainage area of Canoe Creek, 62 square miles																					
7.0	2.7	1.5	.....	0.06	20.9	0	0	0.05	.....	217	0	.....	18	19.7	198	225	2.9	0.65	.....	4	
RIVER NEAR SICAMOUS																					
0.8	0.7	0.8	.....	.....	2.3	0	.....	.....	.....	26.4 (17.1)	0 (0)	.....	4.5	0	20.0	28.8	6.7	.....	1.8	5	
RIVER NEAR MALAKWA																					
0.7	1.3	1.2	0.12	Trace	10.0	0	0.35	0.15	.....	21.9	0	.....	5.2	2.9	20.9	38.9	9.4	.....	2.0	6	
RIVER (PERRY RIVER) NEAR CRAIGELLACHIE																					
0.6	0.5	0.5	0.64	0.06	3.5	0	0.4	0.05	.....	17.1 (17.1)	0 (0)	.....	4.8	1.0 (0)	15.0 (14.0)	23.8	6.5	.....	2.2	7	
MOUTH, NEAR SPENCES BRIDGE																					
8.6	7.8	1.8	.....	0.04	19.6	0	0	0.05	.....	112 (110)	9.6 (8.4)	.....	4.9	4.2 (6.0)	112 (110)	138	12.9	0.6	.....	8	

TABLE II—Continued

Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection		Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>4</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
				On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
STATION No. 50: NICOLA RIVER AT																			
1	July	29/49....	47	.....	.....	66	.....	..... (0)	8.4 (8.8)	20 (20)	3 .....	.....	.....	228	163	0.222	.....	67.2	40.4
STATION No. 51: COLDWATER																			
2	July	29/49....	81	.....	.....	59	.....	..... (5.0)	7.5 (7.5)	0 (5)	4 .....	.....	.....	113	80.6	0.111	.....	7.8	17.0
STATION No. 52: BONAPARTE																			
3	Aug.	8/50....	160	.....	.....	62	.....	..... (0)	8.2 (8.2)	10 .....	40 (35)	79	71	494	295	0.402	.....	32.4	53.8
STATION No. 53: DEADMAN																			
4	Aug.	8/50....	78	.....	.....	58	.....	..... (0)	8.5 (8.5)	5 (5)	0.5 .....	.....	.....	384	225	0.306	.....	25.0	39.0
STATION No. 54: SETON																			
5	*Aug.	18/50....	124	.....	.....	60	.....	..... (1.5)	7.6 (7.7)	2 ( $<5$ )	3 .....	.....	.....	105	63.8	0.087	.....	9.6	20.8
* Sample partly lost by spillage during transit.																			
STATION No. 55: BRIDGE RIVER (NORTH FORK)																			
6	Apr.	4/51....	55	.....	.....	44	.....	.....	7.7	.....	50	45	40	146	90.6	0.123	.....	29.0	17.3
STATION No. 56: SAN JOSÉ RIVER																			
7	Aug.	19/50**..	123	.....	.....	64	.....	..... (9.4)	8.5 (8.7)	5 (15)	0.9 .....	.....	.....	475	301	0.410	.....	77.2	28.0
** Calcium precipitated, but corrected for loss.																			
STATION No. 57: WILLIAMS LAKE																			
8	Aug.	21/50....	51	.....	.....	69	.....	..... (0)	8.5 (8.8)	10 (30)	0.6 .....	.....	.....	525	345	0.470	.....	91.8	31.3



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalis		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>2</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total <sup>b</sup>			+	-	-	
HIGHWAY BRIDGE NEAR NICOLA																					
8.9	8.0	2.8	.....	0.10	22.4	0 (0)	0.7	0	.....	167 (112)	5.8 (9.6)	8.4	4.0	0	137	175	11.0	0.7	.....	1	
RIVER NEAR MERRITT																					
3.4	2.9	0.6	0.08	0.04	6.3	0.1	Trace	0.05	.....	67.1 (69.5)	0 (0)	8.2	10.2	1.4	56.4	73.6	10.0	.....	1.0	2	
RIVER AT CACHE CREEK																					
22.5	17.4	2.8	4.0	0.05	47.2	0	0.7	0.20	.....	245 (253)	11.5 (2.4)	.....	18	7.0 (11.0)	227 (232)	295	14.3	0.8	.....	3	
RIVER NEAR SAVONA																					
15.5	14.8	2.5	0.10	.....	19.9	0	0.4	0	.....	199 (229)	14.4 (12.2)	20.6	19.8	0	161	224	16.4	0.9	.....	4	
LAKE NEAR LILLOOET																					
1.9	1.8	1.1	.....	0.1	9.9	0 (0)	0	.....	.....	61.0 (42.7)	0 (0)	8.4	6.1	9.7	59.7	71.7	5.9	.....	0.8	5	
NEAR LILLOOET—Drainage area near Bridge River, 1,350 square miles																					
6.4	3.5	1.3	1.8	0.09	13.6	1.6	0	0.10	.....	74.9	0	.....	7.0	8.1	69.5	87.8	9.7	.....	0.7	6	
(LAC LA HACHE) NEAR WRIGHT																					
32.5	26.8	4.8	.....	0.07	10.9	0	1.2	0.15	.....	298 (266)	16.6 (26.4)	16	18	0	204	286	21.0	0.9	.....	7	
AT WILLIAMS LAKE																					
37.5	32.5	4.5	.....	0.08	15.1	0	0.7	0.25	.....	314 (300)	27.4 (30)	.....	11	0 (0)	232 (250)	315	22.9	1.0	.....	8	

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
1	Feb. 19/50....	18			33			7.7	3	5	9.6	5.2	108	71.4	0.097		7.8	10.2
2	Mar. 19.....	17			32.5			7.9	3	5	6.2	5.2	134	80.8	0.110		8.4	11.3
3	Apr. 21.....	18			38			7.9	20	5	14	11	173	112	0.1525		20.4	13.0
4	May 22.....	45			42			7.9	15	5			138					11.1
5	June 21.....	33			53			7.7	5	130	154	145	81.6	62.8	0.085		5.0	9.2
6	July 19.....	86			58			7.8	5	25	28	26	74.7	52.8	0.072		16.8	10.8
7	Aug. 17.....	63			57			7.7	7	7			82.6					10.3
8	Aug. 20*.....	110			60			6.9 (9.7)	3 (1.0)	15 (7.7)			73.9					10.8
9	Sept. 23.....	83			56			7.7	4	25			79.7					9.0
10	Oct. 27.....	87			40			7.8	10	6	5.6	5.3	102	67.4	0.092		8.6	9.3
11	Nov. 23.....	67			33			7.6	10	15			87.8					10.2
12	Dec. 19.....	50			33			7.3	7	5			107					10.7
13	Jan. 20/51....	24			32.5			7.6	5	3			126	81.8	0.111		13.0	11.7
14	Average..... (12 samples)	48			42.3			7.6	8	20			107.8	75.6	0.103		11.4	10.6

\* Field sample, not included in average.

## STATION No. 59: CHILKO RIVER NEAR

			Gauge height in feet																
15	Feb. 19/50....	18	610	602	32.5			7.7	0	5	6.6	4.2	76.8	52.4	0.071	0.086	5.2	10.6	
16	Mar. 19.....	17	640	659	32.5			7.7	0	5	7.2	6.2	86.7	50.0	0.068	0.086	4.0	10.7	
17	Apr. 21.....	18	665	624	36.5			7.8	10	5	7.4	5.4	89.7	58.8	0.080	0.105	5.8	11.8	
18	May 22.....	45	2,470	1,940	45			7.9	5	7			86.0					11.2	
19	June 21.....	49	16,700	9,760	50			7.6	5	140	137	132	61.9	48.6	0.066	2.19	8.0	8.9	
20	July 19.....	86	9,630	10,300	56			7.4	3	25	24	21	59.2	44.8	0.061	1.161	12.0	9.7	
21	Aug. 17.....	63	6,070	6,930	57			7.4	10	15			65.2					9.4	
22	Sept. 23.....	83	4,950	3,490	55			7.7	5	25			55.9					8.3	
23	Oct. 27.....	98	2,050	2,610	40			7.6	7	6	5.4	3.7	63.7	46.2	0.063	0.256	6.2	9.0	
24	Nov. 23.....	67	1,940	2,180	33			7.4	7	7			63.0					10.2	
25	Dec. 19.....	56	1,060	1,210	33			7.4	7	5			66.7					9.2	
26	Jan. 20/51....	37	754	839	32			7.6	5	6	3.8	2.9	80.1	49.6	0.0625	0.101	4.6	11.3	
27	Average..... (12 samples)	53	3,962	3,429	41.9			7.6	5	20			71.2	50.1	0.068		5.5	10.1	

TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index			No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-		

## ALEXIS CREEK—Drainage area, 2,400 square miles

4.4	4.8	1.4	0.55	0.09	9.2	0	0	0.05	.....	61.5	0	10.4	10.6	0	43.5	.....	.....	.....	.....	.....	1
6.1	6.0	1.7	0.66	0.03	16.5	0	0	0.05	0	74.4	0	12.6	11.9	0	53.3	.....	.....	.....	.....	.....	2
8.1	10.2	3.2	0.9	0.09	6.7	0	0	0.10	.....	105	0	18.4	15.7	0	65.7	.....	.....	.....	.....	.....	3
5.8	7.5	2.5	6.5	.....	4.7	0	.....	.....	0.01	83.4	0	.....	16.8	0	51.6	.....	.....	.....	.....	.....	4
2.3	3.3	1.5	5.8	0.33	6.4	0	0	.....	.....	43.4	0	.....	7.4	0	32.5	.....	.....	.....	.....	.....	5
1.9	2.8	1.0	1.2	0.20	4.8	0	0.40	.....	.....	42.0	0	.....	6.2	0.3	34.7	.....	.....	.....	.....	.....	6
2.0	3.2	1.0	.....	.....	8.2	0	.....	.....	0	45.9	0	.....	7.1	0	33.9	.....	.....	.....	.....	.....	7
1.5	2.8	0.9	.....	.....	8.2	0	0	.....	.....	35.4 (32.9)	0 (0)	.....	1.0	4.2 (13.5)	33.2 (50.5)	.....	.....	.....	.....	.....	8
2.3	3.3	1.2	.....	.....	7.4	0	.....	.....	.....	42.7	0	.....	5.0	0	32.0	.....	.....	.....	.....	.....	9
4.3	4.8	1.5	0.6	0.10	4.6	0	0	0.10	.....	61.2	0	.....	9.8	0	40.9	.....	.....	.....	.....	.....	10
3.0	3.5	1.3	.....	.....	2.5	0	.....	.....	0	58.6	0	.....	6.4	0	38.0	.....	.....	.....	.....	.....	11
4.4	4.8	1.5	.....	.....	7.4	0	.....	.....	.....	63.4	0	.....	9.2	0	45.0	.....	.....	.....	.....	.....	12
5.4	5.3	1.4	.....	0.17	3.8	0	0.4	0.05	.....	76.1	0	12.8	10.8	0	51.4	.....	.....	.....	.....	.....	13
4.2	4.9	1.6	.....	0.14	6.9	0	0.1	0.07	.....	63.1	0	.....	9.3	0	43.7	68.9	18.9	.....	.....	1.0	14

## REDSTONE—Drainage area, 3,230 square miles

1.9	2.2	0.7	0.13	0.09	10.5	0	0	0	0.11	39.3	0	5.2	4.8	2.0	34.2	.....	.....	.....	.....	.....	15
2.4	2.4	0.8	0.05	0.03	17.3	0	0	0	.....	42.7	0	4.4	4.6	0.2	36.6	.....	.....	.....	.....	.....	16
2.1	3.2	1.0	0.5	0.08	10.5	0	0	0.08	.....	47.6	0	6.2	5.8	0	38.0	.....	.....	.....	.....	.....	17
1.5	3.2	0.9	.....	.....	8.4	0	.....	.....	0.05	46.4	0	.....	8.8	0	34.1	.....	.....	.....	.....	.....	18
1.2	1.6	0.8	8.2	0.32	6.4	0	0	0	.....	30.7	0	.....	4.7	1.9	27.1	.....	.....	.....	.....	.....	19
0.8	1.6	0.8	1.4	0.32	6.4	0	0.4	.....	.....	31.7	0	.....	4.2	1.5	27.5	.....	.....	.....	.....	.....	20
0.6	1.7	0.6	.....	.....	6.6	0	.....	.....	0	29.8	0	.....	4.6	1.6	26.0	.....	.....	.....	.....	.....	21
0.8	1.5	0.7	.....	.....	5.6	0	.....	.....	.....	28.1	0	.....	3.1	1.0	24.0	.....	.....	.....	.....	.....	22
1.3	1.8	0.7	0.5	0.12	5.6	0	0.4	0.10	.....	33.4	0	.....	4.4	0.6	28.0	.....	.....	.....	.....	.....	23
1.4	1.8	0.9	.....	.....	4.5	0	.....	.....	0.02	41.5	0	.....	3.1	0	31.0	.....	.....	.....	.....	.....	24
1.5	1.6	0.5	.....	.....	6.7	0	.....	.....	.....	34.2	0	.....	4.6	1.1	29.1	.....	.....	.....	.....	.....	25
1.7	2.1	0.6	0.23	0.04	3.9	0	0.4	0	0	43.9	0	4.8	3.1	0	35.2	.....	.....	.....	.....	.....	26
1.4	2.1	0.8	.....	0.14	7.7	0	0.2	0.03	.....	37.4	0	.....	4.6	0.2	31.0	49.5	12.5	.....	.....	1.2	27



TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>6</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 60: HORSEFLY RIVER AT

1	Aug. 19/50....	123	341	411	64	.....	.....	7.9 (8.0)	5 (10)	0.9 (clear)	.....	.....	94.7	61.0	0.083	0.056	8.8	14.0
---	----------------	-----	-----	-----	----	-------	-------	--------------	-----------	----------------	-------	-------	------	------	-------	-------	-----	------

STATION No. 61: QUESNEL RIVER AT GRAVELLE FERRY,

			Gauge height in feet															
2	Feb. 19/50....	13	2,820	3,000	35	.....	.....	8.1	5	5	11.2	9.0	138	90.0	0.1225	0.684	12.4	23.2
3	Mar. 19.....	17	2,390	2,440	35	.....	.....	8.0	0	0.3	.....	.....	151	89.0	0.121	0.574	6.8	23.3
4	Apr. 19.....	20	3,440	3,150	38	.....	.....	7.9	25	55	66	63	158	103	0.140	0.956	9.8	24.2
5	May 20.....	10	10,400	9,740	42	.....	.....	8.2	20	25	.....	.....	139	.....	.....	.....	.....	21.6
6	June 19.....	24	29,700	23,100	55	.....	.....	7.9	5	200	258	245	121	83.2	0.113	6.67	7.2	20.5
7	July 19.....	86	16,800	18,400	61	.....	.....	7.9	3	50	59	55	113	77.4	0.105	3.51	9.2	20.4
8	Aug. 19.....	61	8,360	9,390	58	.....	.....	7.8	7	9	.....	.....	115	.....	.....	.....	.....	20.7
9	Aug. 22*.....	106	8,080	9,390	63	.....	.....	7.9 (9.9)	3 (1.0)	15 (7.9)	.....	.....	116	.....	.....	.....	.....	19.0
10	Sept. 24.....	82	5,430	7,750	58	.....	.....	8.0	2	15	.....	.....	117	.....	.....	.....	.....	19.4
11	Oct. 19.....	94	5,590	5,520	46	.....	.....	7.7	4	10	18	17	117	77.2	0.105	1.16	6.4	19.4
12	Nov. 19.....	71	5,100	5,660	34	.....	.....	7.6	5	8	.....	.....	122	.....	.....	.....	.....	20.5
13	Dec. 19.....	50	4,110	4,320	34	.....	.....	7.6	6	5	.....	.....	174	.....	.....	.....	.....	28.8
14	Jan. 19/51....	25	3,630	3,730	32	.....	.....	7.8	0	3	.....	.....	138	88.8	0.121	0.870	9.2	23.8
15	Average..... (12 samples)	46	8,148	8,017	44	.....	.....	7.9	7	25	.....	.....	134	86.9	0.118	.....	8.7	22.1

\* Field sample, not included in average.

STATION No. 62: COTTONWOOD

16	Aug. 23/50....	133	.....	.....	64	.....	.....	7.7 (9.7)	15 (1.0)	6 (7.8)	14 (20)	12 (5)	107	72.8	0.099	.....	19.2	15.6
----	----------------	-----	-------	-------	----	-------	-------	--------------	-------------	------------	------------	-----------	-----	------	-------	-------	------	------

TABLE II  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation Index		No.
	Sodium (Na)	Potassium (K)	Total	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-	

HIGHWAY BRIDGE AT HORSEFLY

2.5	1.8	0.9	.....	0.09	5.4	0 (0)	1.1	0	.....	48.8 (40.2)	0 (2.4)	8.2	8.7	5.2 (6.0)	45.2 (44.0)	58.4	7.6	.....	0.8	1
-----	-----	-----	-------	------	-----	----------	-----	---	-------	----------------	------------	-----	-----	--------------	----------------	------	-----	-------	-----	---

NEAR QUESNEL—Drainage area, 3,900 square miles

3.2	1.1	0.3	0.28	0.08	9.6	0	0.6	0	0.06	73.2	1.2	5.0	4.2	9.1	71.1	.....	.....	.....	.....	2
3.8	1.8	0.5	.....	0.05	15.2	0	0.5	0	.....	81.3	0	4.8	4.8	6.1	73.7	.....	.....	.....	.....	3
3.7	2.6	0.7	3.7	0.23	10.7	0	0.6	0.10	.....	87.1	0	7.2	6.4	4.2	75.6	.....	.....	.....	.....	4
3.4	1.3	0.4	.....	.....	9.3	0	.....	0	.....	70.0	3.6	.....	4.5	4.5	67.9	.....	.....	.....	.....	5
1.6	1.4	0.6	14.7	0.23	7.9	0	0.9	0.05	.....	62.5	2.4	.....	5.0	2.5	57.7	.....	.....	.....	.....	6
2.4	1.6	0.4	2.2	0.09	7.1	0	0.8	0	.....	65.0	0	.....	6.2	9.2	60.8	.....	.....	.....	.....	7
1.6	0.9	0.3	.....	.....	11.1	0	0	.....	0	58.6	0	.....	3.8	10.2	58.2	.....	.....	.....	.....	8
2.3	1.1	0.3	.....	.....	5.3 (8.0)	0 (0)	.....	.....	.....	66.1 (61.7)	0 (0)	.....	3.8	2.7 (8.4)	56.9 (59.0)	.....	.....	.....	.....	9
2.2	1.7	0.4	.....	.....	10.1	0	.....	.....	.....	64.7	0	.....	3.3	4.5	57.5	.....	.....	.....	.....	10
2.7	1.1	0.3	1.4	0.05	7.7	0	0.4	0.05	.....	68.3	0	.....	6.4	3.5	50.5	.....	.....	.....	.....	11
2.9	1.3	0.4	.....	.....	6.8	0	.....	.....	0.01	68.1	0	.....	4.3	7.2	63.0	.....	.....	.....	.....	12
4.5	2.2	0.8	.....	.....	10.7	0	.....	.....	.....	100	0	.....	2.4	8.3	90.5	.....	.....	.....	.....	13
3.2	1.0	0.2	.....	0.27	7.6	0	0.4	0.05	.....	78.1	0	.....	5.8	3.8	72.5	.....	.....	.....	.....	14
2.9	1.5	0.4	.....	0.14	9.5	0	0.6	0.04	.....	72.9	0.6	.....	4.6	7.4	67.1	78.3	4.5	.....	0.4	15

RIVER NEAR QUESNEL

3.2	1.8	0.4	0.70	0.06	8.6	0	0	0.10	.....	59.8 (58.6)	0 (0)	.....	7.0 (6.0)	3.1	52.1	65.6	6.9	.....	0.9	16
-----	-----	-----	------	------	-----	---	---	------	-------	----------------	----------	-------	--------------	-----	------	------	-----	-------	-----	----

TABLE II—Continued

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>4</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 63: NECHAKO RIVER

			Gauge height in feet															
			†	†														
1	Apr. 17/50....	15			33			7.0	20	9	13	11	224	114	0.155		10.6	27.2
2	May—No sample taken.....			14,800														
3	June 18.....	39	27,700	26,600	65			7.6	20	5	19	16	88.5	59.6	0.081	4.46	14.8	11.9
4	July—No sample taken.....			29,900														
5	Aug. 18.....	78	16,500	17,400	63			7.4	15	0.4			76.4	55.8	0.076	2.49	18.4	9.4
6	Sept. 15.....	70	11,000	11,000				7.7	10	2			81.8					10.3

† Records at Isle Pierre, drainage area 16,200 square miles.

STATION No. 64: CHILAKO RIVER NEAR

7	Aug. 25/50....	112			61			7.9 (9.2)	20 (3.0)	6 (8.1)			284 (25)					30.0 (5)
---	----------------	-----	--	--	----	--	--	--------------	-------------	------------	--	--	-------------	--	--	--	--	-------------

STATION No. 65: NECHAKO RIVER AT

8	Sept. 18/50....	110			62			6.8 (10.6)	20 (1.5)	5 (7.5)			55.9 (10)					10.8 (clear)
---	-----------------	-----	--	--	----	--	--	---------------	-------------	------------	--	--	--------------	--	--	--	--	-----------------

STATION No. 66: STUART RIVER AT FORT

9	Mar. 21/50....	15	1,430*	1,480	34			7.9	0	0.4			106	69.8	0.095	0.269	16.0	13.2
10	Apr. 15.....	12	1,180*	1,250	39			7.9	8	2			99.2	76.0	0.103	0.242	19.0	15.6
11	May 16.....	14	3,080	3,200	34			7.7	7	0.5			80.5					10.9
12	June 15.....	27	7,630*	7,740	60			7.8	5	2			93.5	127	0.173	2.620	54.6	13.2
13	July 15.....	58	10,700*	10,200	61			7.7	20	0.5			92.4	74.2	0.101	2.144	14.2	13.2
14	Aug. 15.....	65	7,260*	7,090	58			8.0	20	3			90.2					13.1
15	Aug. 25*.....	143	6,030	7,090	67			7.8 (9.8)	8 (0)	0.8 (8.2)			89.6	63.6	0.087	0.403	16.4	12.8 (20)
16	Sept. 14.....	167	4,320*	4,260				7.6	20	1			90.1					12.4
17	Oct. 17.....	49	2,590*	2,620	43			7.8	3	2			92.4	67.0	0.091	0.468	22.4	11.9

\*—estimated

\* Field sample, not included in average.



TABLE II—Continued

## Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non-car- bonate	Total <sup>b</sup>			+	-	

## NEAR MOUTH AT PRINCE GEORGE

5.7	2.0	0.7	1.1	0.10	11.7	0	0.7	0.10	.....	99.8	0	8.4	5.9	9.6	91.4	103	4.5	.....	1.1	1
2.9	2.4	0.9	0.8	0.04	4.6	0	1.3	0.10	.....	49.8	0	.....	5.4	0.8	41.6	54.0	10.5	.....	1.1	2
2.2	2.0	0.4	.....	0.12	3.0	0	0.4	.....	0.001	41.5	0	.....	5.0	0	32.6	42.9	11.6	.....	1.5	3
2.4	1.8	0.4	.....	.....	3.1	.....	.....	.....	.....	48.8	0	.....	4.9	0	35.6	46.9	9.8	.....	1.1	4
																				5
																				6

## MOUTH, NEAR PRINCE GEORGE

14.5	6.8	2.5	.....	.....	6.0	0 (0)	.....	.....	.....	184 (181)	0 (0)	.....	11.2	0	135	162	9.7	0.05	.....	7
------	-----	-----	-------	-------	-----	----------	-------	-------	-------	--------------	----------	-------	------	---	-----	-----	-----	------	-------	---

## HIGHWAY BRIDGE AT VANDERHOOF

3.1	2.0	0.3	.....	.....	15.2	0	0	.....	.....	34.2 (25.6)	0 (0)	.....	2.4	7.2	35.2	50.6	9.8	.....	2.2	8
-----	-----	-----	-------	-------	------	---	---	-------	-------	----------------	----------	-------	-----	-----	------	------	-----	-------	-----	---

## ST. JAMES—Drainage area, 5,400 square miles

3.5	1.8	0.4	.....	0.05	12.3	0	0	0.10	.....	57.1	0	5.8	4.9	0	47.3	.....	.....	.....	.....	9
3.6	2.0	0.5	.....	0.10	16.6	0	0	0.15	0	53.7	0	8.0	5.4	9.7	53.7	.....	.....	.....	.....	10
2.3	1.7	0.3	.....	.....	5.4	0	.....	.....	.....	46.8	0	.....	4.6	0	36.7	.....	.....	.....	.....	11
2.9	2.0	0.4	.....	0.04	6.1	0	0	0.10	.....	53.4	0	.....	4.8	1.0	44.8	.....	.....	.....	.....	12
3.2	2.2	0.4	.....	0.11	4.9	0	0	0.10	.....	52.5	0	9.6	6.1	3.1	46.1	.....	.....	.....	.....	13
2.3	2.0	0.3	.....	.....	8.8	0	.....	.....	0.01	51.2	0	.....	5.7	0.2	42.2	.....	.....	.....	.....	14
2.7	1.9	0.6	.....	0.09	4.3	0	0.4	0.12	.....	54.2 (50.0)	(0)	.....	4.8	5.0	0	43.0	.....	.....	.....	15
3.2	2.0	0.6	.....	.....	4.7	0	.....	.....	.....	51.2	0	.....	6.0	2.0	44.0	.....	.....	.....	.....	16
2.8	1.8	0.4	.....	0.07	5.6	0	0.4	0	.....	53.2	0	4.2	5.4	0	41.2	.....	.....	.....	.....	17

TABLE II—Continued  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Continued  
(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge <sup>a</sup> (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance K x 10 <sup>4</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		

STATION No. 66: STUART RIVER AT

			Gauge height in feet															
1	Nov. 14.....	58	1,760 <sup>e</sup>	1,760	33			7.6	10	4			91.6					13.5
2	Dec. 14.....	6	1,460 <sup>e</sup>	1,460	34			7.8	15	2			96.2					15.2
3	Jan. 16/51....	28	1,220	1,220	32			7.5	20	0.4			95.3					13.1
4	Feb. 15.....	33	1,120 <sup>e</sup>	1,110	32			7.7	20	0.3			107	68.2	0.093	0.206	19.2	14.1
5	Average..... (12 samples)	43	3,646	3,615	38			7.8	12	1.5			94.5	80.4	0.109		24.2	13.2

<sup>e</sup>—estimated

STATION No. 67: NECHAKO RIVER AT FORT

6	Feb. 15/50....	17		8,900	36			7.3	5	2			55.8	42.6	0.058		12.0	8.8
7	Mar. 16.....	11			33			7.7	7	3			58.7	41.6	0.0566		7.2	8.2
8	Apr. 15.....	12			33			7.5	5	3			49.7	40.6	0.055		10.0	7.6
9	May 15/50....	15	6,570	7,810	48			7.5	25	6			66.4					9.6
10	June 15.....	27	15,500	15,740	62			7.3	5	5	12.2	6.4	50.0	80.6	0.110	3.37	47.6	6.9
11	July 15.....	58	16,900	16,260	62			7.2	10	0.9			45.0	35.6	0.048	1.62	8.6	6.4
12	Aug. 15.....	65	8,250	8,340	61			7.4	10	3			46.2					8.5
13	Sept. 16.....	165	5,310	5,350	60			7.4	10	0.6			45.8					6.3
14	Oct. 14.....	52	4,220	4,360	45			7.6	2	2			46.4	36.6	0.050	0.417	10.0	6.7
15	Nov. 14.....	76	4,370	4,310	35			7.5	7	4			35.7					6.1
16	Dec. 16.....	48		ice	33			7.5	7	15			50.3					8.5
17	Jan. 16/51....	28		conditions	32			7.2	10	3			47.4	39.8	0.054		11.8	6.4
18	Average..... (12 samples)	48	8,303	8,884	45			7.4	9	4			49.8	45.3	0.0616		15.3	7.5

STATION No. 68: NORTHERLY

19	Aug. 26/50....	60			60			7.8 (10.4)	10 (0)	0.6 (8.2) (20)			88.2	66.4	0.090		16.4	13.6
----	----------------	----	--	--	----	--	--	---------------	-----------	----------------------	--	--	------	------	-------	--	------	------

STATION No. 69: STELLAKO

20	Sept. 1/50....	136			61			7.8 (10.2)	6 (2.5)	1 (7.8)			90.7	71.4	0.097		23.0	11.6
----	----------------	-----	--	--	----	--	--	---------------	------------	------------	--	--	------	------	-------	--	------	------

(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Borax (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		
	Sodium (Na)	Potassium (K)	Total <sup>a</sup>	Dissolved								Gravi- metric	Colori- metric	Non- sur- bonate	Total <sup>b</sup>			+	-	No

## FORT ST. JAMES—Continued

4.0	2.0	0.6			0	0	0.03	53.2	3.5		3.2	0.1	50.1				1
3.5	1.8	0.3			0	0		56.1	0		4.8	6.1	52.1				2
3.1	0.5	0.3		4.6	0			52.5	0		5.0	2.5	45.5				3
3.2	1.7	0.4	0.04	5.8	0	0.7	0	58.3	0	7.6	4.7	0.5	48.3				4
3.1	1.8	0.4	0.07	6.2	0	0.1	0.06	53.2	0		5.5	2.1	45.7	56.7	7.8	0.8	5

FRASER—Drainage area, 6,700 square miles

0.9	1.3	0.3	.....	0.06	3.5	0	0	0.05	0.16	31.7	0	3.6	4.4	0	26.7	.....	.....	.....	.....	6
1.5	1.8	0.4	.....	0.04	5.4	0	Trace	0.05	.....	34.2	0	6.2	4.2	0	26.7	.....	.....	.....	.....	7
1.5	1.2	0.5	.....	0.10	11.4	0	0	0.10	.....	24.4	0	4.2	3.4	5.2	25.2	.....	.....	.....	.....	8
1.6	2.0	0.7	.....	.....	8.4	0	.....	.....	0	41.0	0	.....	5.8	0	30.5	.....	.....	.....	.....	9
1.1	1.6	0.3	0.6	0.08	3.8	0	0	0.05	.....	27.3	0	.....	4.2	0	21.7	.....	.....	.....	.....	10
0.8	1.3	0.3	.....	0.08	3.6	0	0	0	.....	22.0	0	5.8	3.1	1.3	19.3	.....	.....	.....	.....	11
0.8	1.5	0.2	.....	.....	7.4	0	.....	.....	0.01	24.4	0	.....	4.6	4.5	24.5	.....	.....	.....	.....	12
1.2	1.7	0.4	.....	.....	3.1	0	.....	.....	.....	26.8	0	.....	5.1	0	20.5	.....	.....	.....	.....	13
0.9	1.4	0.2	.....	0.07	5.9	0	0.4	0.10	.....	24.4	0	4.2	4.6	0.4	20.4	.....	.....	.....	.....	14
1.3	1.5	0.5	.....	.....	2.5	0	.....	.....	0	25.6	0	.....	4.4	0	20.5	.....	.....	.....	.....	15
1.4	1.2	0.3	.....	.....	6.8	0	.....	.....	.....	29.3	0	.....	3.8	3.0	27.0	.....	.....	.....	.....	16
1.2	1.2	0.7	.....	0.12	3.0	0	0.4	0.10	.....	26.8	0	5.2	4.7	0	20.9	.....	.....	.....	.....	17
1.2	1.5	0.4	.....	0.8	5.4	0	.....	0.06	0.07	28.1	0	.....	4.4	0.6	23.6	34.4	11.9	.....	1.7	18

## RIVER NEAR FORT FRASER

2.3	3.3	0.7	.....	0.05	4.3	0 (0)	0.4	0.15	.....	56.1 (46.4)	0 (2.4)	6.4	4.0	0 (2.0)	43.4 (44.0)	57.2	13.0	.....	0.05	10
-----	-----	-----	-------	------	-----	----------	-----	------	-------	----------------	------------	-----	-----	------------	----------------	------	------	-------	------	----

## RIVER NEAR FORT FRASER

2.3	3.9	1.0	.....	0.16	4.3	0	0.7	.....	56.6 (50.0)	..... (0)	7.8	5.2	0	38.4	57.1	17.6	.....	1.0	20
-----	-----	-----	-------	------	-----	---	-----	-------	----------------	--------------	-----	-----	---	------	------	------	-------	-----	----



TABLE II—Concluded

**Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Concluded**

(In parts per million)

No.	Date of collection	Storage period (Days)	Stream discharge* (Second-feet)		Water temperature (°F.)	Dissolved oxygen	Carbon dioxide	pH	Colour	Turbidity	Suspended matter		Specific conductance  K x 10 <sup>3</sup> at 25°C.	Residue on Evaporation dried at 105°C. (Dissolved solids)			Loss on ignition at 550°C.	Calcium (Ca)
			On sampling date	Monthly mean							Dried at 105°C.	Ignited at 550°C.		P.P.M.	Tons per acre-foot	Thousand tons per day		
STATION No. 70: BURNS LAKE																		
1	Sept. 1/50....	136	.....	.....	59 (9.2)	..... (4.0)	..... (7.5)	35 (60)	3	.....	.....	108	101	0.1375	.....	44.6	14.1	
STATION No. 71: WILLOW																		
2	Aug. 23/50....	133	.....	.....	66 (9.2)	..... (1.5)	9.7 (7.9)	25 (25)	6 (10)	7.6	5.2	104	72.0	0.098	.....	17.8	15.0	
DATA SUPPLIED BY																		
3	Nov. 1944—Thompson River at Aschroft, B.C.....							7.7	.....	.....	.....	.....	78.7	.....	.....	.....	.....	
4	Aug. 1940—Mountain stream near Chase, B.C.....							8.0	.....	.....	.....	.....	371.8	.....	.....	.....	.....	
5	Nov. 1944—Coquitlam River near Port Coquitlam, B.C.....							7.2	.....	.....	.....	.....	42.9	.....	.....	.....	.....	
6	Nov. 1944—Lytton Creek at Lytton, B.C.....							6.9	.....	.....	.....	.....	121.6	.....	.....	.....	.....	
7	Nov. 1944—Hallecks Creek at North Bend, B.C.....							3.0	.....	.....	.....	.....	78.7	.....	.....	.....	.....	
8	Nov. 1944—Murray and Waterfall Creeks near Spences Bridge, B.C.....							8.4	.....	.....	.....	.....	200.2	.....	.....	.....	.....	

TABLE II—Concluded  
Chemical Analyses of Surface Waters in the Fraser River Drainage Basin—Concluded  
(In parts per million)

Magnesium (Mg)	Alkalies		Iron (Fe)		Sulphate (SO <sub>4</sub> )	Chloride (Cl)	Nitrate (NO <sub>3</sub> )	Fluoride (F)	Boron (B)	Bicarbonate (HCO <sub>3</sub> )	Carbonate (CO <sub>3</sub> )	Silica (SiO <sub>2</sub> )		Hardness as CaCO <sub>3</sub>		Sum of Constituents	Per cent sodium	Saturation index		No.	
	Sodium (Na)	Potassium (K)	Totale	Dissolved								Gravi- metric	Colori- metric	Non- car- bonate	Total <sup>b</sup>			+	-		
NEAR BURNS LAKE																					
3.5	4.1	1.6	.....	0.30	1.6	0	2.7	.....	.....	58.6 (51.0)	0 (0)	9.4	7.2	0	49.6	70.8	14.7	.....	1.1	1	
RIVER AT WILLOW RIVER																					
2.7	1.8	0.4	0.72	0.11	6.7	0	0.4	.....	.....	61.0	0	.....	7.2	0	48.5	64.3	7.4	.....	0.9	2	
CANADIAN PACIFIC RAILWAYS																					
.....	.....	.....	.....	.....	8.7	5.2	.....	.....	.....	48.8	0	1.4	.....	2.9	42.9	.....	.....	.....	.....	3	
.....	.....	.....	.....	.....	8.7	2.6	.....	.....	.....	139.6	0	14.3	.....	0	114.4	.....	.....	.....	.....	4	
.....	.....	.....	.....	.....	5.8	0.9	.....	.....	.....	17.4	0	10.0	.....	2.9	17.2	.....	.....	.....	.....	5	
.....	.....	.....	.....	.....	2.9	7.8	.....	.....	.....	87.2	0	10.0	.....	5.7	77.2	.....	.....	.....	.....	6	
.....	.....	.....	.....	.....	7.7	5.2	.....	.....	.....	61.1	0	10.0	.....	0	42.9	.....	.....	.....	.....	7	
.....	.....	.....	.....	.....	9.7	3.5	.....	.....	.....	189.8	5.7	10.0	.....	0	160.2	.....	.....	.....	.....	8	

## PART II

### MUNICIPAL WATERS WITHIN THE FRASER RIVER DRAINAGE BASIN

When in 1949 survey studies were being carried out in the Columbia River basin a number of municipal water supplies within the lower Fraser River drainage basin were also studied. The remainder were sampled and field-tested in 1950 when almost the entire accessible portion of the basin was travelled with the mobile laboratory. At that time information on the operation of many of the civic water systems was also obtained. Information on others was obtained by co-operation of municipal officials or taken from the Regional Industrial Indices of British Columbia<sup>1</sup>.

Much of the available information on all these systems is condensed below under the headings: population, ownership, source, treatment, storage capacity, water consumption and industrial use.

The chemical quality of the civic water sampled is shown in Table III. Sum of constituents and saturation index have also been reported for each water in Table III.

Appendix B lists the incorporated municipalities and other communities which are known to have organized water systems; their locations are shown in Figure 2 (in pocket) in such a manner that the water hardness of the supplies, when known, is also classified.

Table IV is a summary of the information available regarding basin area studied, total basin population and the population served with water by organized systems.

In Table V the available information on waterworks systems in the basin, such as source, treatment and hardness of the water supply is summarized.

### DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS

#### GREATER VANCOUVER WATER DISTRICT

Within the Fraser River delta, in an area about 20 miles wide and extending roughly 60 to 70 miles upriver, a number of incorporated cities and district municipalities have formed a Water District governed by a Board on which each member community is represented. This District Board, which includes waterworks engineers and other trained personnel, owns and operates the supply of water to the entire district but most municipalities own and operate their own distribution system. Some of the municipalities are also served by other systems and from other water sources.

In 1948, metropolitan Vancouver was considered to include 5 cities, 10 district municipalities\* and 2 unorganized districts totalling 563 square miles in area. By 1950 it was expected that 370 square miles of this area would be served by the Greater Vancouver Water District<sup>2</sup>.

This Water District is still expanding as population and industrial development in the delta area grows and other sources of supply become inadequate. In 1947, *population served* was estimated at 494,100 and in 1949 the Water District reported a population of 610,817. Comparison of the latter figure with population figures based on the 1951 census and with other data indicates close agreement in all cases except Vancouver which is some 77,840 lower by census calculation. Assuming the figures based on the ninth census to be more correct, then about 529,000 persons were served by this District in 1950-51. It is however difficult to determine exact figures owing to rapid expansion within the area and the fact that several municipalities are served with water by other sources than those of the Water District.

The *sources of supply* for the Water District are from rivers and lakes in the coastal range lying north of the Fraser River and Burrard Inlet. This protected catchment area of, eventually, some 225.7 square miles is either owned or leased by the District Board.

The main sources are:

1. Coquitlam River or Lake—capacity 400 m.g.d.
2. Seymour River with headwaters in Loch Lomond and Burwell Lake—capacity 220 m.g.d.
3. Capilano River with headwaters in Palisades Lake—capacity 200 m.g.d.

These waters flow by gravity to the various systems with no treatment, although chlorination is available on all supplies, if considered necessary owing to work being carried out in the watersheds.

\* Incorporated district municipalities in British Columbia are somewhat similar to townships in Eastern Canada and may have within their jurisdiction several relatively large communities.

<sup>1</sup> Regional Index of British Columbia—Regional Development Division, Department of Trade and Industry of British Columbia, 1949 edition and 1952 edition.

<sup>2</sup> Water Supply to Rural Areas around Vancouver—W. H. Powell, Water and Sewage, (now Municipal Utilities) June, 1948, p. 21.



Water consumption in 1948 (371 day-period) by the Water District according to the administrative Board was as follows:

South of Burrard Inlet:

From Capilano River.....	10,236·670	million gallons		
From Seymour River.....	7,844·455	“	“	
				18,081·125 million gallons

North of Burrard Inlet:

From Capilano and Seymour Rivers...	405·731	m.g.		
Coquitlam River.....	2,066·877	m.g.		
			2,472·608	“ “
Total.....			20,553·733	“ “

or a daily average of 55·401 m.g.

The north shore area<sup>1</sup> which is included in that portion north of Burrard Inlet is only partially served by the Greater Vancouver Water District. The portion served used in 1948, 476,858 m.g. or 1·285 m.g.d. Except for this north shore portion there is a per capita consumption of 105 gallons within the Water District. Presuming a similar per capita consumption in the north shore area then about 12,240 persons are served within this area.

*Storage capacity* for the Water District is considerable because the headwaters of the systems are protected lakes and rainfall on the watershed is heavy. (See Vancouver, B.C. page 70.)

*Industrial use* within the Water District is varied and quite high. (See below under individual communities served.)

Municipalities served in all or part by the Greater Vancouver Water District in 1950 were Vancouver, New Westminster, Port Coquitlam, Port Moody and North Vancouver, District Municipalities of North Vancouver, West Vancouver, Coquitlam, Maple Ridge, Pitt Meadows, Burnaby, Delta, Surrey, Fraser Mills and Richmond. Two unorganized areas were also served.

<sup>1</sup> Presumably the portions of North Vancouver City, North Vancouver District Municipality and West Vancouver District Municipality served.

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
(Fraser River Drainage Basin)  
BRITISH COLUMBIA

	<b>ABBOTSFORD</b> —160 acres (Incorporated as a village, Feb. 1924)			<b>ARMSTRONG*</b> —1,140 acres (Incorporated as a city, 1913)			
	1941*	1949	1951*	1941	1949	1950	1951
Municipality.....							
Population served:							
In municipality.....	562	800	785	977	1,100	1,100	1,126
Outside municipality.....	.....	380	.....	.....	500	.....	.....
Total.....	.....	1,180	.....	.....	1,600	.....	.....
Date(s) of survey.....	August 24, 1950.....			July 28, 1949; Feb. 13, 1950.....			
Ownership.....	Municipally owned and operated.....			Municipally owned and operated.....			
Source of supply.....	Springs nearby.....			Davis (Fortune) Creek.....			
Treatment.....	No treatment; water is pumped direct from collecting reservoirs to system.			No treatment; water from dammed creek flows by gravity to reservoir and system.			
Storage capacity (thousand gallons).....	Reservoirs..... 296			1 reservoir..... 60			
Consumption (average in m.g.d.).....	1 tank..... 72			1 reservoir (1950)..... 160			
	No record.....			1948..... 1949.....			
				0.48 (est.)..... 0.48.....			
Industrial use.....	An agricultural and shopping centre; no major industrial user.			Fruit growing and farming; two packing companies, a sawmill and a supplier of compressed oxygen.			
Remarks.....	*Populations according to eighth and ninth census of Canada.			*Armstrong itself lies within the Columbia River drainage basin but civic water is taken from Fraser River system.			

	<b>BRIDGEPORT</b> (Unincorporated)	<b>BRIGHOUSE</b> (Unincorporated)
Municipality.....		
Population served:		
In municipality.....		
Outside municipality.....		
Total.....		
Date(s) of survey.....		
Ownership.....		
Source of supply.....		
Treatment.....	See Richmond District Municipality	
Storage capacity (thousand gallons).....		
Consumption (average in m.g.d.).....		
Industrial use.....		
Remarks.....	A part of and included in data given on Richmond District Municipality.	

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

<b>ASHCROFT</b> (Unincorporated)			<b>BEACH GROVE*</b> (Unincorporated)		<b>BRALORNE</b> (Unincorporated)	
1941	1949	1951	1951		1951	
.....	.....	1,118	.....		1,500	
.....	.....	.....	.....		.....	
.....	850	.....	400†		.....	
August 8, 1950.....			1951.....		April 20, 1951.	
Privately owned and operated by Ashcroft Water, Electric & Improvement Co., Ltd.			Beach Grove Water Board.....		Owned and operated by Bralorne Mines Ltd.	
Thompson River.....			No data; presumed to be wells and springs.....		Blackbird Creek, fed by mountain springs.	
Chlorination; water is pumped from Thompson River to ground reservoir on nearby hill; then flows by gravity to system.			No data.....		No treatment; water flows by gravity to reservoirs and system.	
Underground reservoir..... 135			No data.....		2 tanks.....100 total	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data.....		.....	
.....			No data			

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

Municipality.....	<b>BURQUITLAM</b> (Unincorporated)		<b>CANOE</b> (Unincorporated)	
	1950	1951	1948	1951
Population served:				
In municipality.....	.....	.....	.....	1,100
Outside municipality.....	.....	.....	.....	.....
Total.....	500 (est.)	590 (est.)	900 (est.)	.....
Date(s) of survey.....	August 19, 1950.....		1950.....	
Ownership.....	Municipally owned and operated—Clinton Waterworks District.		Cloverdale Water Co., Ltd.....	
Source of supply.....	A mountain creek.....		Springs and artesian wells.....	
Treatment.....	No treatment; water flows by gravity to system.		No data. (presumably no treatment).....	
Storage capacity (thousand gallons).....	No data.....		No data.....	
Consumption (average in m.g.d.).....	No record (80 services).....		No data.....	
Industrial use.....	No major industrial user.....		No data.....	
Remarks.....			Cloverdale lies within the Surrey District Municipality.	



**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
(Fraser River Drainage Basin)  
BRITISH COLUMBIA

<b>CHASE</b> (Unincorporated)		<b>CHILLIWACK</b> —1,040 acres (Incorporated as a city, Feb. 1908)			<b>CHILLIWACK DISTRICT MUNICIPALITY*</b> —53,000 acres (Incorporated)		
1949	1951	1941	1949	1951	1941	1949	1951
.....	.....	3,675	5,000	5,663	7,787	.....	.....
.....	.....	.....	6,000*	.....	.....	.....	.....
400*	700*	.....	11,000	.....	.....	6,000	11,000
1950.....		August 24, 1949.....			August 24, 1949..		
Privately owned by Carlin Bros. Ltd.; Chase Waterworks District.		Owned and operated by the Elk Creek Waterworks Co., Ltd., Chilliwack.					
No data.....		Elk, Dunville and Nevin Creeks, rising in hills nearby. Famihi and Lihumitson Creeks are also available if required.					
No data.....		Chlorination begun late in 1949; all three creeks are screened at intakes into small wooden reservoir and the water then flows by gravity to system. Major supply is Elk Creek (2 m.g.d.)			See Chilliwack		
No data.....		Open reservoirs—125, 200 and 100; construction underway to increase reservoir capacity at intakes to 2 m.g. total.					
No data.....		1948					
No data.....		4.0(est.)†					
No data.....		Main users are a cannery (peas, vegetables, fruit) and a frozen food plant. Area is a centre of extensive agricultural and fruit-growing activity.					
*Estimated from number of services.....		*Chilliwack District Municipality.....			*Includes communities of Rosedale, Sardis, Vedder Crossing and Yarrow but not all served with water. Total population in 1951—13,592.		
		† Includes pumpage into District Municipality.					
<b>COQUITLAM DISTRICT MUNICIPALITY</b> —37,204 acres* (Incorporated)			<b>CRESCENT BEACH</b> (Unincorporated)		<b>DELTA DISTRICT MUNICIPALITY*</b> —80,818 acres (Incorporated, November, 1879)		
1941	1949	1951	1951		1941	1949	1951
7,949	.....	15,697	.....		4,287	.....	6,700
.....	.....	.....	.....		.....	.....	.....
.....	13,000	.....	400*		.....	5,000	.....
.....			March, 1951.....		August 8, 1949.....		
.....			Privately owned and operated by Crescent Beach Waterworks Ltd.		Delta Municipal Waterworks. North Delta Waterworks. Beach Grove Water Board. Tswassen Water Board.		
Supplied by Greater Vancouver Water District.			Two artesian wells.....		Springs and wells and from Greater Vancouver Water District.		
See Vancouver, B.C.....			No treatment; water is pumped from wells to reservoir and system.		See Vancouver and Greater Vancouver Water District, Beach Grove, etc.		
No data.....			1 tank..... 60		Reservoir..... 200		
No data.....			No record (390 services).....		1 tank..... 18		
No data.....			No major industrial user; tourism.....		1948-49		
					0.85		
					Main activity in area is agriculture, peat-processing and fishing.		
* Includes communities of Mallardville, Burquitlam and Essondale; 23,000 acres served by Greater Vancouver Water District.			Crescent Beach lies within the Surrey District Municipality.		* Includes the communities of Beach Grove, Kennedy, Sunbury and Ladner; only 39,889 acres served by Greater Vancouver Water District.		
			* Being a seaside resort, population rises to 2,500 in summer. System is expanding rapidly.				

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

	<b>EAST RICHMOND</b> (Unincorporated)	<b>EBURNE</b> (Unincorporated)	<b>ENDERBY—655 acres</b> (Incorporated as a city, 1905)		
			1941	1949	1951
Municipality.....					
Population served:					
In municipality.....			538	.....	888
Outside municipality.....			.....	.....	.....
Total.....			.....	1,000	.....
Date(s) of survey.....			August 4, 1950.....		
Ownership.....			Municipally owned and operated.....		
Source of supply.....	A part of Richmond District Municipality and included in data given on this Municipality.		Brash Creek and a well (23 ft. deep) alongside Shuswap River.		
Treatment.....			No treatment: Brash Creek flows by gravity after settling in wooden tanks to system. The well water, said to be river water naturally filtered, is pumped direct to system.		
Storage capacity (thousand gallons).....			Open reservoir..... 300		
Consumption (average in m.g.d.).....			1949-50 0-25 (est.)		
Industrial use.....			Main users are C.P.R., a sawmill and a creamery.		
Remarks.....					

	<b>HANEY*</b> (Unincorporated)		<b>HARRISON HOT SPRINGS</b> 1,678 acres (Incorporated as a village, May, 1949)	
	1949	1951	1949	1951
Municipality.....				
Population served:				
In municipality.....		2,700		477
Outside municipality.....		.....		.....
Total.....	2,300	.....	470	.....
Date(s) of survey.....	December, 1949.....		August, 1950.....	
Ownership.....	Municipally owned and operated.....		No organized system.....	
Source of supply.....	Supplied by Greater Vancouver Water District.		Privately owned wells, mountain creek and Harrison Lake.	
Treatment.....	See Vancouver and Greater Vancouver Water Board.			
Storage capacity (thousand gallons).....	None.....			
Consumption (average in m.g.d.).....	No data.....			
Industrial use.....	Mill work (sash and doors) and other smaller industries.		A summer resort with hot mineral springs.....	
Remarks.....	* Haney lies within and is included in data given on Maple Ridge District Municipality.			

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

<b>ESSONDALE</b> (Unincorporated)			<b>FRASER MILLS DISTRICT MUNICIPALITY—390 acres</b> (Incorporated)			<b>GOLD BRIDGE</b> (Unincorporated)		
1949	1951		1941	1949	1951	1949	1951	
.....	.....		552	.....	370	140 (est.)	150 (est.)	
.....	.....		.....	.....	.....	.....	.....	
3,395	.....		.....	450 (est.)	.....	.....	.....	
December, 1949.....			December, 1949.....			1948-49.....		
.....			Privately owned and operated by the Canadian Western Lumber Co., Ltd.			Gold Bridge Waterworks District.		
.....			Supplied by Greater Vancouver Water District.			No data.		
Essondale is a part of and included in data given on Coquitlam District Municipality.			See Vancouver and Greater Vancouver Water District.			No data.		
None.....			No data.....			No data.		
No data.....			No record: included in data given for Greater Vancouver Water District.			No record (38 services).		
No data.....			Lumbering.....			No data.		
Essondale is a provincial institution within Coquitlam District Municipality but it buys direct from Greater Vancouver Water District.								

<b>HOPE—1,200 acres</b> (Incorporated as a village, 1929)			<b>IOCO</b> (Unincorporated)			<b>KAMLOOPS—912 acres</b> (Incorporated as a city, July, 1893)		
1941	1950	1951	1949	1951		1941	1949	1951
515	.....	1,668	.....	.....		5,959	.....	8,099
.....	.....	.....	.....	.....		.....	.....	.....
.....	3,000	.....	350 (est.)	400 (est.)		.....	10,500*	.....
August 17, 1950.....			1949.....			July 29, 1949.....		
Municipally owned and operated.....			Owned and operated by Imperial Oil Ltd.....			Municipally owned and operated.		
Schkam, Pringle and Charles Creeks, principally Schkam Creek.			No data.....			South Thompson River.		
No treatment; waters flow by gravity finally from Schkam Creek to reservoir in hills and to system.			No data.....			Chlorination: turbine-pumped from sump well at river to reservoirs and system.		
Open natural reservoir..... 500			No data.....			1 closed reservoir..... 1,500		
No record (480 services).....			1949.....			1 tank..... 150		
			No data			1948-49.....		
			(93 services)			1.5†		
						(Maximum—4.0)		
Two sawmills; C.P.R. has its own water system.			Oil refining and oil storage.....			Irrigation (hop-growing, etc.) may account for 50% of total. C.P.R. shops are a large user; 3 canneries, a brewery and lumbering.		
.....			.....			* Probably includes North Kamloops village and district.		
.....			.....			† Total area consumption.		

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

Municipality.....	<b>KENNEDY</b> (Unincorporated)	<b>LADNER</b> (Unincorporated)	<b>LILLOOET</b> —152 acres (Incorporated as a village, Dec. 1946)		
			1941	1950	1951
Population served:		1951			
In municipality.....		2,000	.....	.....	469
Outside municipality.....		.....	.....	.....	.....
Total.....		.....	.....	350*	.....
Date(s) of survey.....			August 18, 1950.....		
Ownership.....			Privately owned and operated by Lillooet Waterworks District.		
Source of supply.....			Springs and creeks†.....		
Treatment.....	Included in data on Delta District Municipality.		No treatment; springs and creeks in mountains flow to concrete reservoir then by gravity to Lillooet District Waterworks system. There is a small settling basin at the intake to the concrete reservoir.		
Storage capacity (thousand gallons).....			Concrete reservoir.....		50
Consumption (average in m.g.d.).....			1949-50		
			No record (95 services)		
Industrial use.....			Irrigation is the main use. A cannery uses the railway well and Cayoos Creek. Lillooet is a divisional point on the Pacific Great Eastern Railway.		
Remarks.....			* Total population estimated at 600. † Also one area of town has no organized supply while another area is served by gravity from a mountain creek.		

Municipality.....	<b>MATSQUI DISTRICT MUNICIPALITY</b> 54,165 acres (Incorporated)		<b>McBRIDE</b> —240 acres (Incorporated as a village, 1932)		
	1941	1951	1941	1949	1951
Population served:					
In municipality.....	5,601*	10,308*	237	435 (est.)	490
Outside municipality.....	.....	.....	.....	.....	.....
Total.....	.....	.....	.....	.....	.....
Date(s) of survey.....	April 6, 1951 (by questionnaire).....		1951.....		
Ownership.....	Privately owned and operated by Matsqui Water Co., Ltd.		Owned and operated by C.N.R.....		
Source of supply.....	A mountain lake.....		Dominion Creek.....		
Treatment.....	No treatment; water enters system by gravity.		No data.....		
Storage capacity (thousand gallons).....	No data.....		No data.....		
Consumption (average in m.g.d.).....	No data.....		No data (148 services).....		
Industrial use.....	No major industrial user.....		A divisional point on the C.N.R. and a lumbering centre.		
Remarks.....	* Not all served; system is limited and only a few farms in area are served (200 services).—estimated population served 900.				



# DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS

## (Fraser River Drainage Basin)

### BRITISH COLUMBIA

LYTTON (Incorporated as a village)			MAILLARDVILLE (Unincorporated)			MAPLE RIDGE DISTRICT MUNICIPALITY—66,000 acres (Incorporated)		
1941	1950	1951				1941	1949	1951
.....	.....	312	.....	.....	.....	6,476	.....	9,891
.....	.....	.....	.....	.....	.....	.....	.....	.....
.....	600*	.....	.....	.....	.....	.....	10,000*	.....
August 9, 1950.....						December, 1949.		
Lytton Water Supply Company†.....						Municipally owned and operated.		
Lytton Creek.....			A part of and included in data given for Coquitlam District Municipality.			Supplied by Greater Vancouver Water District.		
No treatment; water flows by gravity direct to system from tank in nearby hills.						See Vancouver and Greater Vancouver Water District.		
1 tank.....						None.		
No record (100 services).....						1949	1950	
						0-46	0-55	
C.P.R.....						Some of the main activities in district are small fruit, dairy, poultry and fur farming.		
* Probably not all served.....						* Not all served.		
† C.P.R. maintains the water mains into town in return for water rights for railway use.								

MERRITT—1,679 acres (Incorporated as a city, April, 1911)			MISSION CITY—821 acres (Incorporated as a village, Dec. 1922)			MISSION DISTRICT MUNICIPALITY 73,000 acres (Incorporated)		
1941	1949	1951	1941	1949	1951	1941	1949	1951
940	.....	1,251	1,957	3,090	2,668	2,718	.....	4,449*
.....	.....	.....	.....	810*	.....	.....	.....	.....
.....	1,500*	.....	.....	3,900	.....	.....	810†	.....
July 29, 1949.....			August 23, 1949.....			August 23, 1949.		
Municipally owned and operated.....			Municipally owned and operated.....					
Coldwater River.....			Cannell Lake, Cedar Valley springs and Silverdale Creek; principally Cannell Lake.			A portion of District is supplied by Mission City Waterworks.		
Chlorination (sodium hypochlorite solution); water enters underground filtration gallery and is then pumped to reservoir and system.			No treatment; water flows by gravity from all sources to system. The springs and lake supply the higher portion of town.			See Mission City.		
1 elevated tank..... 250			Open reservoir at springs..... 175			See Mission City.		
			Open dam reservoir at creek..... 1,000					
			Cannell Lake..... 22,324,000 acre-ft.					
			1948			1948		
						0-103		
	1948							
	0-25 (est.)							
Main activities in area are cattle raising and lumbering. C.P.R. also uses civic water.			Total..... 0-489			See Mission City.		
			This is an important retail and wholesale distributing centre; canning, fruit and milk processing, sawmilling and manufacture of matches.					
* Total population—2,000.....			* Probably in Mission District Municipality...			* Not all served.		
						† Based on data supplied by Mission City Waterworks' officials in 1949.		

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

Municipality.....	<b>NEWTON STATION</b> (Unincorporated)	<b>NEW WESTMINSTER</b> —4,394 acres (Incorporated as a city, July, 1860)												
Population served:		<table> <tr> <th>1941</th><th>1949</th><th>1951</th></tr> <tr> <td>21,967</td><td>.....</td><td>27,789</td></tr> <tr> <td>.....</td><td>.....</td><td>850</td></tr> <tr> <td>.....</td><td>34,000</td><td>28,639</td></tr> </table>	1941	1949	1951	21,967	.....	27,789	.....	.....	850	.....	34,000	28,639
1941	1949	1951												
21,967	.....	27,789												
.....	.....	850												
.....	34,000	28,639												
In municipality.....														
Outside municipality.....														
Total.....														
Date(s) of survey.....		December, 1949.....												
Ownership.....		Municipally owned and operated.....												
Source of supply.....	A part of and included in data given for Surrey District Municipality.	Supplied by the Greater Vancouver Water District mostly from Seymour River and Coquitlam Lake.												
Treatment.....		See Vancouver and Greater Vancouver Water District.												
Storage capacity (thousand gallons).....		1 reservoir..... 6,900 1 reservoir..... 2,300												
Consumption (average in m.g.d.).....		1949 5.0												
Industrial use.....		This is a distributing centre for products of Fraser River valley and has diversified industries including salmon-canning, lumbering, machinery manufacture and meat-packing.												
Remarks.....														

Municipality.....	<b>PITT MEADOWS DISTRICT MUNICIPALITY</b> —12,000 acres (Incorporated)	<b>PORT MANN</b> (Unincorporated)												
Population served:	<table> <tr> <th>1941</th><th>1949</th><th>1951</th></tr> <tr> <td>1,119</td><td>.....</td><td>1,434</td></tr> <tr> <td>.....</td><td>.....</td><td>.....</td></tr> <tr> <td>.....</td><td>840</td><td>.....</td></tr> </table>	1941	1949	1951	1,119	.....	1,434	.....	.....	.....	.....	840	.....	
1941	1949	1951												
1,119	.....	1,434												
.....	.....	.....												
.....	840	.....												
In municipality.....														
Outside municipality.....														
Total.....														
Date(s) of survey.....	1949-50.....													
Ownership.....	Municipally owned and operated.....													
Source of supply.....	Supplied by Greater Vancouver Water District, mostly from Coquitlam Lake.	Included in and part of Surrey District Municipality.												
Treatment.....	See Vancouver and Greater Vancouver Water District.													
Storage capacity (thousand gallons).....	None.....													
Consumption (average in m.g.d.).....	1949 0.14													
Industrial use.....	Area activity is primarily agriculture and lumbering; some manufacturing.													
Remarks.....														

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

<b>NORTH KAMLOOPS</b> (Incorporated)			<b>NORTH VANCOUVER</b> —3,131 acres (Incorporated as a city, March, 1906)			<b>NORTH VANCOUVER DISTRICT MUNICIPALITY*</b> —40,818 acres† (Incorporated)		
1941	1949	1951	1941	1949	1951	1941	1949	1951
.....	.....	1,979	8,914	.....	15,687	5,931	.....	14,469
.....	.....	.....	.....	.....	.....	.....	.....	.....
.....	2,000	.....	.....	11,000	.....	.....	11,000	.....
July 29, 1949.....			August 9, 1949.....			1949.....		
Municipally owned and operated.....			Municipally owned and operated.....			Municipally owned and operated.....		
Supplied by Kamloops Municipal Waterworks...			Lynn Creek, 4 miles distant; also from Greater Vancouver Water District.			Lynn, Mosquito and Kilmer Creeks; Kennedy Lake: supplied also by Greater Vancouver Water District.		
See Kamloops, B.C.....			Lynn Creek is used directly with no treatment. See also Greater Vancouver Water District and Vancouver.			No treatment except chlorination on certain creek waters: waters enter system by gravity: some booster pumping.		
			Rice Lake.....30,000			No data.....		
			1948-49.....1950.....			1949.....1950.....		
			3.0.....3.5			0.28.....0.38		
			Fishing and lumbering with a substantial ship building industry; a highly industrialized area.			A highly industrialized area. See North Vancouver.		
						* Includes communities of Capilano, Caulfield, Deep Cove, Dollarton and Lynn Creek.		
						† 11,300 acres served by Greater Vancouver Water District.		

<b>PORT COQUITLAM</b> —6,200 acres (Incorporated as a city, March 1913)			<b>PORT MOODY</b> —2,980 acres* (Incorporated as a city, March, 1913)			<b>PRINCE GEORGE</b> —1,092 acres (Incorporated as a city, March, 1915)		
1941	1949	1951	1941	1949	1951	1941	1950	1951
1,539	.....	3,232	1,512	.....	2,246	2,027	4,500*	4,703
.....	.....	.....	.....	.....	.....	.....	.....	.....
.....	2,500	.....	.....	2,500	.....	.....	.....	.....
August 9, 1949.....			August 9, 1949.....			August 24, 1950.....		
Municipally owned and operated.....			Municipally owned and operated.....			Municipally owned and operated.....		
Supplied by Greater Vancouver Water District.			Supplied by Greater Vancouver Water District.			Nechako River.		
See Vancouver and Greater Vancouver Water District.			See Vancouver and Greater Vancouver Water District.			Chlorination and natural filtration, from river above town through 300 ft. of gravel into 20 ft. well then pumped to reservoir on hill, or river water may be pumped direct. Standby source is gravel well, 20 ft. from river.		
None.....			1 reservoir.....1,000			1 elevated tank (open).....100		
						A 3 m.g. open reservoir is planned.		
						1949-50.....		
						0.65 (est.)		
0.50 (approx.)			0.75 (approx.)			Main users are sawmills, a dairy, and a roofing company. Prince George is a main distributing centre.		
Freight terminus of C.P.R.; lumbering; rubber manufacturing and fruit farming.			Main activities are sawmilling, lumbering and oil-refining.			* Total district population—10,000.		
			* Only about 1,980 acres served by Greater Vancouver Water District.					

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

	<b>VANCOUVER</b> —27,965 acres (Incorporated as a city, April, 1886)			<b>VEDDER CROSSING</b> (Unincorporated)
	1941	1949	1951	
Municipality.....				No data.....
Population served:				
In municipality.....	275,353	421,471	342,728	
Outside municipality.....	.....	1,200*	2,105*	
Total.....	.....	422,671†	344,833	
Date(s) of survey.....	August 9, 1949.....			
Ownership.....	Municipally owned and operated.....			A part of Chilliwack District Municipality.
Source of supply.....	Supplied by the Greater Vancouver Water District from Capilano River, Seymour Lake and Coquitlam Lake. The city is supplied mainly from the first two sources.			A portion of population including a federal defence installation are supplied with water from Chilliwack River which is not treated except for chlorination. See Table II, Station No. 22.
Treatment.....	No treatment; water from behind a dam on the Capilano River at Palisades Lake is piped, after screening, by gravity to system. Similarly at dams at Burwell Lake and Loch Lomond, Seymour River water is piped by gravity to system. Coquitlam Lake is also piped by gravity to various portions of Greater Vancouver Water District system.			
Storage capacity (thousand gallons).....	Tank..... 750 Palisade Lake (Capilano River)....16,000 acre ft. Loch Lomond, Seymour River, (Burwell Lake)..... 5,400 acre ft. Coquitlam Lake.....12,000 acre ft.			
Consumption (average in m.g.d.).....	Total capacity.....33,400 acre ft. 1948 43.0 (approx.)			
Industrial use.....	Many types of industry located in this area; but many of the larger are outside the city proper.			
Remarks.....	Chlorination available on all supplies if required. * University Endowment Lands. † Data supplied by Greater Vancouver Water District.			



**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

WALHACHIN (Unincorporated)	WELLS (Unincorporated)	WEST VANCOUVER DISTRICT MUNICIPALITY—20,515 acres.* (Incorporated)
1948	1951	1941      1949      1951
.....	.....	7,669      .....      13,990
.....	.....	.....
60 (est.)	1,000	.....      11,000      .....
1948-49.....	January, 1951.....	August 9, 1949.
Walhachin Water District.....	Privately owned, and operated by the Cariboo Gold Quartz Mining Co., Ltd.; (Wells Townsite Co., Ltd.)	Municipally owned and operated.
No data.....	Mosquito Creek and Red Gulch Creek. Jack of Clubs Lake is used for fire protection if necessary.	Supplied from Greater Vancouver Water District. Only partly served.
No data.....	No treatment; creek waters enter reservoirs and system by gravity.	See Vancouver and Greater Vancouver Water District.
No data.....	2 tanks in town, each..... 6 1 tank at the mine..... 30	2 tanks, each..... 100 Eagle Lake..... 33,000
No data (13 services).....	1950	1949
	Domestic..... 0.030	1.5
	Industrial..... 0.025	
	Total..... 0.055	
No data.....	Only industrial user is the Cariboo Gold Quartz Mining Co., Ltd.	Mainly a residential community.
		* In 1950 about 7,680 acres served by Greater Vancouver Water District.

**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

	<b>WHALLEY</b> (Unincorporated)	<b>WHITE ROCK*</b> (Unincorporated)		
		1949	Population rises in summer to 10,000-15,000. Week-end summer population may rise to 40,000.	
Municipality.....		6,000†		
Population served:				
In municipality.....				
Outside municipality.....				
Total.....				
Date(s) of survey.....		March 1, 1950.....		
Ownership.....	A part of	Privately owned and operated by White Rock Waterworks Co., Ltd.		
Source of supply.....	and included in	Graeine spring and three flowing wells.		
Treatment.....	Surrey	No treatment; water is pumped to reservoirs and system. Civic supply is mixture of various sources.		
	District			
Storage capacity (thousand gallons).....	Municipality	3 elevated tanks..... 20, 50 and 50		
		2 ground reservoirs..... 42 and 120		
		Total..... 282		
Consumption (average in m.g.d.).....		1949		
		0.22		
		(Maximum—0.68)		
Industrial use.....		Mainly a tourist resort.....		
Remarks.....		* A part of Surrey District Municipality.		
		† A bulk supply of water is also supplied to 300 services in the Sunnyside area of the District.		

Municipality.....	<b>WILLIAMS LAKE</b> —324 acres (Incorporated as a village, March 1929)		
	1941	1950	1951
Population served:			
In municipality.....	540	.....	913
Outside municipality.....	.....	.....	.....
Total.....	.....	1,200	.....
Date(s) of survey.....	August 21, 1950.		
Ownership.....	Municipally owned and operated.		
Source of supply.....	Williams Lake.		
Treatment.....	Chlorination (sodium hypochlorite); water is pumped from near shore to reservoir and system.		
Storage capacity (thousand gallons).....	1 elevated tank..... 85		
	Pacific Great Eastern Railway..... 60		
Consumption (average in m.g.d.).....	1949		
	0.80		
Industrial use.....	Main user is Pacific Great Eastern Railway.		
Remarks.....	Main area activity is cattle raising.		



**DESCRIPTION OF MUNICIPAL WATERWORKS SYSTEMS**  
**(Fraser River Drainage Basin)**  
**BRITISH COLUMBIA**

	<b>WHALLEY</b> (Unincorporated)	<b>WHITE ROCK*</b> (Unincorporated)
Municipality.....		<div>1949</div> <div>Population rises in summer to 10,000-15,000. Week-end summer population may rise to 40,000.</div>
Population served:		
In municipality.....		6,000†
Outside municipality.....		.....
Total.....		.....
Date(s) of survey.....		March 1, 1950.....
Ownership.....	A part of	Privately owned and operated by White Rock Waterworks Co., Ltd.
Source of supply.....	and included in	Grasine spring and three flowing wells.
Treatment.....	Surrey	No treatment; water is pumped to reservoirs and system. Civic supply is mixture of various sources.
	District	
Storage capacity (thousand gallons).....	Municipality	<div>3 elevated tanks..... 20, 50 and 50</div> <div>2 ground reservoirs..... 42 and 120</div> <div>Total..... 282</div>
Consumption (average in m.g.d.).....		<div>1949</div> <div>0.22</div> <div>(Maximum—0.68)</div>
Industrial use.....		Mainly a tourist resort.....
Remarks.....		* A part of Surrey District Municipality. † A bulk supply of water is also supplied to 300 services in the Sunnyside area of the District.

	<b>WILLIAMS LAKE—324 acres</b> (Incorporated as a village, March 1929)
Municipality.....	<div>1941</div> <div>1950</div> <div>1951</div>
Population served:	
In municipality.....	540
Outside municipality.....	.....
Total.....	.....
Date(s) of survey.....	August 21, 1950.
Ownership.....	Municipally owned and operated.
Source of supply.....	Williams Lake.
Treatment.....	Chlorination (sodium hypochlorite); water is pumped from near shore to reservoir and system.
Storage capacity (thousand gallons).....	<div>1 elevated tank..... 85</div> <div>Pacific Great Eastern Railway..... 60</div>
Consumption (average in m.g.d.).....	<div>1949</div> <div>0.80</div>
Industrial use.....	Main user is Pacific Great Eastern Railway.
Remarks.....	Main area activity is cattle raising.





TABLE III  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

	Municipality.....	ABBOTSFORD	ARMSTRONG	ASHCROFT	BEACH GROVE	BRALORNE	BRIDGEPORT	BRIGHOUSE	BURKEVILLE
No.	Source(s).....	Springs	Fortune Creek	Thompson River	Springs and wells	Blackbird Creek	Supplied by Richmond District Municipality		
		Raw and finished water	Raw and finished water	Raw and finished water		Raw and finished water			
	Sampling point.....	Town tap	Town tap	Direct from river					
1	Laboratory number.....	3520	3489	4619			See Richmond District Municipality.		
2	Field number.....	505	466	706					
3	Date of collection.....	Aug. 24/49	July 28/49	Aug. 8/50					
4	Storage period (days).....	68	82	84					
5	Sampling temperature, °C.....	17.0		16.7					
6	Test temperature, °C.....	20.5	22.0	26.2					
7	Dissolved oxygen.....								
8	Carbon dioxide (CO <sub>2</sub> ).....	(1.2)	(3.0)	(2.0)					
9	pH.....	7.6 (7.5)	8.1 (8.3)	7.8 (8.0)					
10	Colour.....	5 (5)	0 (5)	8 (6)					
11	Turbidity.....	2	0.5	5.5 (5)					
12	Suspended matter, dried at 105°C.....								
13	Suspended matter, ignited at 550°C.....								
14	Residue on evaporation, dried at 105°C.....	91.8	124		No data; presumed to be similar to wells at Crescent Beach and White Rock, B.C.	No data.			
15	Ignition loss at 550°C.....	32.8	6.4						
16	Specific conductance (micromhos at 25°C.).....	107	180	86.5					
17	Calcium (Ca).....	12.5	32.6	11.5					
18	Magnesium (Mg).....	3.0	2.3	1.6					
19	Iron (Fe) Total.....								
20	Dissolved.....	0.09	0.08						
21	Sodium (Na).....	3.7	1.6	1.8					
22	Potassium (K).....	0.4	1.6	0.8					
23	Carbonate (CO <sub>3</sub> ).....	0 (0)	0 (0)	0					
24	Bicarbonate (HCO <sub>3</sub> ).....	43.9 (43.9)	97.6 (97.6)	41.7					
25	Sulphate (SO <sub>4</sub> ).....	5.4	16.5	5.8					
26	Chloride (Cl).....	5.8 (6.0)	0 (0)	0					
27	Fluoride (F).....	0	0				0.1 p.p.m.† (1952)		
28	Nitrate (NO <sub>3</sub> ).....	15.9	Trace						
29	Silica (SiO <sub>2</sub> ) Gravimetric.....	15	8.8						
30	Colorimetric.....	17	11.2	4.2					
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	36.0	80.0	34.2 (32.0)					
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	7.6	10.8	1.1 (5.0)					
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....	43.6	90.8	35.3 (37.0)					
34	Sum of Constituents.....	85.4	114	46.2					
35	Saturation index.....	-1.2	+0.06	-0.9					
	Remarks:	Note rather high nitrate content.			Beach Grove lies within Delta District Municipality.	†Analysis supplied by Department of National Health and Welfare.	Communities lying in and part of the Richmond District Municipality.		

TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

BURNS LAKE	BURNAST	BURQUITLAM	CANOE	CHILLIWACK			CHILLIWACK DISTRICT MUNICIPALITY	CLINTON	No.
Burns Lake	Supplied by Greater Vancouver Water District		East Canoe Creek; supplied by Salmon Arm Waterworks.	Elk River	Nevin Creek	Dunville Creek	Chilliwack Waterworks	A mountain creek	
Raw and finished water				Raw and finished water				Raw and finished water	
Direct from lake				At intake	At intake	At intake		Town tap	
4813	See Vancouver, B.C.	See Salmon Arm and Salmon Arm District Municipality.		3437	3521	3436	See Chilliwack, B.C.	4567	1
764				510	509	508		728	2
Sept. 1/50				Aug. 25/49	Aug. 25/49	Aug. 25/49		Aug. 19/50	3
136				34	67	34		53	4
15.0				11.0	9.0	9.5		12.2	5
16.8				21.0	20.7	21.0		21.0	6
..... (9.2)									7
..... (4.0)				..... (1.2)		..... (3.0)			8
7.5 (7.5)				8.1 (8.4)	8.1 (8.4)	8.0 (8.4)		8.2 (8.2)	9
35 (60)				3 (5)	5 (5)	3 (5)		3 (15)	10
2.5				0.3	0.7	0.5		0.7 (clear)	11
.....									12
101				154	158	188		236	13
44.6				24.6	28.0	33.8		27.4	14
108				238	243	285		377	15
14.1				41.3	45.8	52.8		51.9	16
3.5				3.0	4.0	3.8		17.5	17
.....				0.04					18
0.30					0.05	0.11		0.08	19
4.1				3.3	2.3	3.7		5.6	20
1.6				0.4	0.3	0.6		1.1	21
0 (0)				2.4 (0)	2.2 (0)	2.8 (0)		4.8 (7.2)	22
71.5 (62.2)				90.3 (95.2)	127 (129)	132 (140)		252 (239)	23
1.6				39.5	28.0	40.2		11.7	24
0				1.6	0.8 (0)	2.1		0 (0)	25
.....				0.05	0	0		0.1	26
2.7				1.2	2.2	3.5		0	27
9.4				6.6	4.8	6.0			28
7.2				10.9	7.4	9.7		20	29
49.6				78.0 (78.0)	104 (106)	113 (115)		202 (208)	30
0				37.3	26.8	34.3		0 (4.0)	31
49.6				115	131	147		202 (212)	32
70.3				148	155	185		237	33
-1.1				+0.1	+0.3	+0.3		+0.7	34
									35

TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

No.	Municipality.....	CLOVERDALE	COQUITLAM DISTRICT MUNICIPALITY	CRESCENT BEACH		DELTA DISTRICT MUNICIPALITY	EAST RICHMOND	EBURNE
	Source(s).....	Springs and artesian wells	Supplied from Greater Vancouver Water District	Artesian wells		Supplied from Greater Vancouver Water Dis- trict and from springs and wells.	Supplied from Greater Vancouver Water District	
				Raw and finished water				
	Sampling Point.....			Well No. 1	Well No. 2			
1	Laboratory number.....			5026	5025			
2	Field number.....			811	810			
3	Date of collection.....			Apr. 30/51	Mar. 17/51			
4	Storage period (days).....			30	44			
5	Sampling temperature, °C.....			4.4	5.6			
6	Test temperature, °C.....			22.8	22.8			
7	Dissolved oxygen.....							
8	Carbon dioxide (CO <sub>2</sub> ).....							
9	pH.....			7.8	8.0			
10	Colour.....			2	0			
11	Turbidity.....			6	0.5			
12	Suspended matter, dried at 105°C.....			1.7				
13	Suspended matter, ignited at 550°C.....			0.9				
14	Residue on evaporation, dried at 105°C.....			125	130			
15	Ignition loss at 550°C.....			22.8	21.8			
16	Specific conductance (micromhos at 25°C.)....	No data.	See Vancouver, B.C.	182	193	See Vancouver, B.C. and Beach Grove, B.C.	See Richmond District Municipality, B.C.	
17	Calcium (Ca).....			16.9	17.8			
18	Magnesium (Mg).....			6.7	7.2			
19	Iron (Fe) Total.....			0.26				
20	Dissolved.....			0.03	0.04			
21	Sodium (Na).....			13.0	10.3			
22	Potassium (K).....			2.0	2.1			
23	Carbonate (CO <sub>3</sub> ).....			0	0			
24	Bicarbonate (HCO <sub>3</sub> ).....			91.0	99.6			
25	Sulphate (SO <sub>4</sub> ).....			7.0	6.0			
26	Chloride (Cl).....			10.3	10.7			
27	Fluoride (F).....	1.2† (1952)		0	0			
28	Nitrate (NO <sub>3</sub> ).....			0	0.4			
29	Silica (SiO <sub>2</sub> ) Gravimetric.....							
30	Colorimetric.....			23	20			
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....			69.7	74.0			
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....			0	0			
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....			69.7	74.0			
34	Sum of Constituents.....			124	124			
35	Saturation index.....			-0.6	-0.3			
	Remarks:	Cloverdale lies with- in Surrey District Mun- icipality, B.C.  †Analyses supplied by Dept. of National Health and Welfare.						



TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

ENDERBY		ESSONDALE	FRASER MILLS DISTRICT MUNICIPALITY	GREATER VANCOUVER WATER DISTRICT	HANEY	HARRISON HOT SPRINGS*	HOPE	
Brash Creek	Well	Supplied from Greater Vancouver Water Dis- trict through Coquitlam District Municipality.	Supplied from Greater Vancouver Water District	Capilano, Coquitlam and Seymour Rivers	Supplied from Greater Vancouver Water Dis- trict through Maple Ridge District Municipality	Mineral hot spring	Schkam Creek	
Raw and finished water				Raw and finished water			Raw and finished water	
Town tap	At pump			Direct from rivers		Direct from spring	Town tap	Town tap
4485	4543	See Vancouver, B.C.	See Vancouver, B.C.	See Vancouver, B.C.	See Vancouver, B.C.	4808	3532	4599
692	694					721	513	723
Aug. 4/50	Aug. 4/50					Aug. 17/50	Aug. 26/49	Aug. 17/50
27	52					151	68	69
15.6						> 60		19.7
25.5 (18.9)	25.5					21.5	19.0	20.5 (22.5)
(3.0)	(40)						(3.0)	(3.0)
7.7 (7.8)	7.4 (7.1)					8.0	7.5 (7.9)	7.2 (7.4)
10 (5)	10					5	5 (5)	3 (<5)
1	25 (25)					0.2	0.4	0
	81							
	78							
90.0	457					1358	48.4	42.6
8.8	82.0					38.6	7.4	5.4
135	725					1965	69.8	67.5
20.4	49.5					83.0	11.6	8.6
2.9	41.9					0	2.3	1.7
	4.25							
0.04	0.10					0.03	0.03	0.05
2.6	50.0					355	1.8	1.2
1.2	2.9					15.0	1.0	0.8
0 (0)	0 (0)					2.4	0 (0)	0 (0)
72.0 (70.8)	383 (381)					17.1	39.6 (41.5)	37.3 (35.4)
9.2	67.7					516	11.9	4.6
0	16.5					283	0	0 (0)
0.15	1.0					3.0	0.05	0
0	2.7					0	0	<0.4
						55	5.0	5.4
6.4	28					64	7.6	7.0
59.0 (58.0)	296 (298)					18.0	32.4 (34.0)	28.5 (29.0)
3.8 (3.0)	0 (0)					189	6.0	0 (0)
62.8 (61.0)	296 (298)					207	38.4	28.5 (29.0)
78.4	449					1330	55.8	42.7
-0.6	+0.1					-0.4	-1.4	-1.8
Fluoride 0.25 p.p.m. (1948); analysis sup- plied by Dept. of National Health and Welfare.	H <sub>2</sub> S present.					*No organized civic supply.		

TABLE III—Continued  
**Chemical Analyses of Civic Water Supplies**  
**FRASER RIVER DRAINAGE BASIN**  
*(In parts per million)*

No.	Municipality.....	KAMLOOPS		KENNEDY	LADNER	LILLOET	LYTTON	MAILLARDVILLE
	Source(s).....	South Thompson River		Supplied from Greater Vancouver Water District through Delta District Municipality		Springs and creeks	Lytton Creek	Supplied from Greater Vancouver Water District through Coquitlam District Municipality
		Raw and finished water				Raw and finished water	Raw and finished water	
	Sampling Point.....	Tap at waterworks plant	Town tap			Town tap	Town tap	
1	Laboratory number.....	3490	4596	See Vancouver, B.C.		4601	4600	See Vancouver, B.C.
2	Field number.....	467	703			726	724	
3	Date of collection.....	July 29/49	Aug. 8/50			Aug. 18/50	Aug. 18/50	
4	Storage period (days).....	81	78			68	68	
5	Sampling temperature, °C.....	19.0	16.7			22.0 (25.3)	15.5	
6	Test temperature, °C.....	22.0	26.2 (25.6)				21.9 (19.3)	
7	Dissolved oxygen.....							
8	Carbon dioxide (CO <sub>2</sub> ).....						(3.0)	
9	pH.....	7.6 (7.9)	7.5 (7.8)			8.1 (8.4)	7.6 (7.8)	
10	Colour.....	7 (6)	10 (5)			2 (<5)	2 (<5)	
11	Turbidity.....	0.2	0.4 (<5)			0.3	0.2	
12	Suspended matter, dried at 105°C.....							
13	Suspended matter, ignited at 550°C.....							
14	Residue on evaporation, dried at 105°C.....	59.4	52.6			216	87.6	
15	Ignition loss at 550°C.....	8.2	8.4			19.8	5.4	
16	Specific conductance (micromhos at 25°C).....	80.5	82.0			347	132	
17	Calcium (Ca).....	12.0	11.6			48.5	22.3	
18	Magnesium (Mg).....	2.1	1.1			19.5	1.6	
19	Iron (Fe) Total.....							
20	Dissolved.....	0.08	0.12			0.04	0.03	
21	Sodium (Na).....	1.3	1.4			5.1	2.6	
22	Potassium (K).....	0.6	0.7			0.4	0.2	
23	Carbonate (CO <sub>3</sub> ).....	0 (0)	0 (0)			9.6 (0)	0 (0)	
24	Bicarbonate (HCO <sub>3</sub> ).....	40.3 (39.1)	40.5 (41.5)			203 (210)	79.3 (78.1)	
25	Sulphate (SO <sub>4</sub> ).....	7.4	7.1			28.0	5.3	
26	Chloride (Cl).....	0 (0)	0			0 (0)	0 (0)	
27	Fluoride (F).....	0.10	0.05			0.05	0	
28	Nitrate (NO <sub>3</sub> ).....	0.35	0			0	0	
29	Silica (SiO <sub>2</sub> ) Gravimetric.....	5.2	6.8			9.6	12	
30	Colorimetric.....	7.6	6.6			9.4	14	
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	33.0 (32.0)	33.2 (34.0)			179 (172)	62.2 (64.0)	
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	5.5	0.2 (1.0)			22.2 (30.0)	0 (1.0)	
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....	38.5	33.4 (35.0)			201 (202)	62.2 (65.0)	
34	Sum of Constituents.....	51.4	48.6			221	84.9	
35	Saturation index.....	-1.2	-1.2			+0.6	-0.7	
	Remarks:	Fluoride—0.07 p.p.m. (1948); analysis supplied by Dept. of National Health and Welfare.						

TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

MAPLE RIDGE DISTRICT MUNICIPALITY	MATSQUI DISTRICT MUNICIPALITY	McBRIDE	MERRITT	MISSION CITY			MISSION DISTRICT MUNICIPALITY	
Supplied from Greater Vancouver Water District	A mountain lake	Dominion Creek	Coldwater River	Cannell Lake	Silverdale Creek	Cedar Valley Springs	Supplied by Mission City Waterworks	
			Raw and finished water	Raw and finished water				
			Town tap	Town tap	At intake	Direct from springs		
See Vancouver, B.C.	No data.	No data.	3491	3530	3528	3529	See Mission City.	1
			469	504	502	503		2
			July 29/49	Aug. 23/49	Aug. 23/49	Aug. 23/49		3
			81	71	71	71		4
			5.0	17.8	13.0	13.5		5
			22.0 (21.8)	19.0	19.5	19.5		6
			(5.0)	(8.5)	(4.0)	(3.9)		7
			7.5 (7.5)	7.1 (6.4)	7.3 (7.3)	7.3 (7.5)		8
			0 (5)	10 (10)	15 (25)	5 (8)		9
			4	0.4	3	0.3		10
			2.0					11
			1.0					12
			80.6	20.6	47.6	73.0		13
			7.8	8.6	13.0	16.0		14
			113	27.7	42.7	76.2		15
			17.0	3.8	7.8	11.2		16
			3.4	0.4	0.8	2.4		17
			0.08					18
			0.04	0.05	0.21	0.03		19
			2.9	1.1	2.4	3.3		20
			0.6	0.1	0.5	0.4		21
			0 (0)	0 (0)	0 (0)	0 (0)		22
			67.1 (69.5)	9.4 (11.0)	24.4 (23.2)	36.6 (39.1)		23
			6.3	6.6	9.1	7.9		24
			0.1 (0)	0 (0)	0	1.5		25
			0.05	0		0.10		26
			Trace	0	0.7	8.0		27
			8.2	3.4	9.4	14		28
			10	4.2	11.8	20		29
			55.0	7.7 (9.0)	20.0 (19.0)	30.0 (32.0)		30
			1.4	3.4	2.8	7.8		31
			56.4	11.1	22.8	37.8		32
			73.6	20.9	45.3	72.8		33
			-1.0	-2.9	-2.0	-1.7		34
								35
		Fluoride 0.10 p.p.m. (1948); analysis sup- plied by Dept. of National Health and Welfare.						

TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

No.	Municipality.....	NEWTON STATION	NEW WESTMINSTER	NORTH KAMLOOPS	NORTH VANCOUVER		NORTH VANCOUVER DISTRICT MUNICIPALITY
	Source(s).....	Supplied from Greater Vancouver Water District through Surrey District Municipality	Supplied from Greater Vancouver Water District	Supplied by Kamloops Municipal Waterworks	Lynn River	Supplied from Greater Vancouver Water District	Supplied from Greater Vancouver Water District and North Vancouver city
					Raw and finished water		
	Sampling Point.....				Direct from river		
1	Laboratory number.....				3439		
2	Field number.....				476		
3	Date of collection.....				Aug. 10/49		
4	Storage period (days).....				49		
5	Sampling temperature, °C.....				17.0		
6	Test temperature, °C.....				21.0 (19.0)		
7	Dissolved oxygen.....						
8	Carbon dioxide (CO <sub>2</sub> ).....						
9	pH.....				7.2 (7.3)		
10	Colour.....				3 (5)		
11	Turbidity.....				0.3 (clear)		
12	Suspended matter, dried at 105°C.....						
13	Suspended matter, ignited at 550°C.....						
14	Residue on evaporation, dried at 105°C.....				18.8		
15	Ignition loss at 550°C.....				6.6		
16	Specific conductance (micromhos at 25°C).....				21.8		
17	Calcium (Ca).....	See Vancouver, B.C.	See Vancouver, B.C.	See Kamloops, B.C.	2.8	See Vancouver, B.C.	See Vancouver, B.C.
18	Magnesium (Mg).....				0.9		and North Vancouver, B.C.
19	Iron (Fe) Total.....						
20	Dissolved.....				0.02		
21	Sodium (Na).....				1.3		
22	Potassium (K).....				0.3		
23	Carbonate (CO <sub>3</sub> ).....				0 (0)		
24	Bicarbonate (HCO <sub>3</sub> ).....				11.6 (7.3)		
25	Sulphate (SO <sub>4</sub> ).....				2.6		
26	Chloride (Cl).....				1.6		
27	Fluoride (F).....				0		
28	Nitrate (NO <sub>3</sub> ).....				0.8		
29	Silica (SiO <sub>2</sub> ) Gravimetric.....						
30	Colorimetric.....				5.8		
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....				9.5 (6.0)		
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....				1.2		
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....				10.7		
34	Sum of Constituents.....				21.8		
35	Saturation index.....				-2.8		
	Remarks:	Newton Station is a part of and included in Surrey District Municipality.			Fluoride—0.05 p.p.m. (1949); analysis supplied by Dept. of National Health and Welfare.		



TABLE III—Continued  
**Chemical Analyses of Civic Water Supplies**  
**FRASER RIVER DRAINAGE BASIN**  
*(In parts per million)*

PITT MEADOWS DISTRICT MUNICIPALITY	PORT MANN	PORT COQUITLAM	PORT MOODY	PRINCE GEORGE		QUESNEL	5 %
Supplied from Greater Vancouver Water District, mostly from Coquitlam River	Included in Surrey District Municipality and supplied from Greater Vancouver Water District	Supplied from Greater Vancouver Water District	Supplied from Greater Vancouver Water District	Nechako River		Quesnel River	
				Raw and finished water		Raw and finished water	
				Plant tap	From sump well, after natural filtration	Town tap	
<i>See Vancouver, B.C.</i>	<i>See Vancouver, B.C.</i>	<i>See Vancouver, B.C.</i>	<i>See Vancouver, B.C.</i>	4570	4569	4785	1
				741	740	737	2
				Aug. 24/50	Aug. 24/50	Aug. 22/50	3
				48	48	134	4
				17.5	14.2	16.2	5
				21.0 (22.2)	16.7 (16.3)	21.0 (21.9)	6
				..... (13.0)	..... (15.0)	..... (7.0)	7
				7.5 (7.2)	7.9 (7.4)	7.5 (7.7)	8
				7 (5)	2 (10)	5	9
				2	2	7	10
				.....	.....	110	11
				.....	.....	8.5	12
				144	259	108	13
				13.6	14.6	38.8	14
				225	382	170	15
				31.2	58.0	27.4	16
				8.1	15.9	3.6	17
				.....	.....	0.16	18
				0.12	0.32	0.03	19
				3.3	4.2	2.8	20
				1.4	2.4	0.6	21
				0 (0)	0 (0)	0	22
				135 (132)	244 (234)	92.7	23
				11.0	25.8	12.7	24
				0 (0)	0 (0)	0 (0)	25
				0.05	0.05	0.10	26
				0	0	2.7	27
				.....	.....	.....	28
				12	27	5.7	29
				111 (108)	200 (192)	76.0	30
				0.1 (14.0)	10.5 (26.0)	7.2	31
				111 (122)	211 (218)	83.2	32
				134	254	101	33
				-0.5	0.4	-0.7	34
				Mn=0.1 p.p.m.	Mn=0.9 p.p.m.	Al=0 p.p.m.	35

TABLE III—Continued  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

No.	Municipality.....	ROSEDALE	RICHMOND DISTRICT MUNICIPALITY	SALMON ARM		SALMON ARM DISTRICT MUNICIPALITY	SARDIS	SOUTH WESTMINSTER
	Source(s).....	Supplied by Chilliwack District Municipality	Supplied from Greater Vancouver Water District	Shuswap Lake	East Canoe Creek	Mostly East Canoe Creek from Salmon Arm Waterworks	Supplied by Chilliwack District Municipality	Supplied by Surrey District Municipality from Greater Vancouver Water District
				Raw and finished water				
	Sampling Point.....			Town tap	From reservoir intake			
1	Laboratory number.....			4566	4565			
2	Field number.....			696	695			
3	Date of collection.....			Aug. 5/50	Aug. 5/50			
4	Storage period (days).....			67	67			
5	Sampling temperature, °C.....			16.5	15.0			
6	Test temperature, °C.....			25.5 (21.7)	25.5 (22.2)			
7	Dissolved oxygen.....							
8	Carbon dioxide (CO <sub>2</sub> ).....			(0)	(3)			
9	pH.....			8.1 (8.5)	8.0 (7.9)			
10	Colour.....			7 (10)	5 (5)			
11	Turbidity.....			6 (8)	0.7			
12	Suspended matter, dried at 105°C.....			7.6				
13	Suspended matter, ignited at 550°C.....			5.2				
14	Residue on evaporation, dried at 105°C.....			82.8	213			
15	Ignition loss at 550°C.....			11.0	9.8			
16	Specific conductance (micromhos at 25°C.)..			128	362			
17	Calcium (Ca).....	See Chilliwack, B.C.	See Vancouver, B.C.	24.6	67.7	See Salmon Arm, B.C.	See Chilliwack, B.C.	See Vancouver, B.C.
18	Magnesium (Mg).....			3.0	7.0			
19	Iron (Fe) Total.....			0.42				
20	Dissolved.....			0.07	0.06			
21	Sodium (Na).....			2.7	2.7			
22	Potassium (K).....			1.1	1.5			
23	Carbonate (CO <sub>3</sub> ).....			0 (4.8)	0 (0)			
24	Bicarbonate (HCO <sub>3</sub> ).....			67.1 (61.1)	217 (210)			
25	Sulphate (SO <sub>4</sub> ).....			9.7	20.9			
26	Chloride (Cl).....			0	0			
27	Fluoride (F).....			0.1	0.05			
28	Nitrate (NO <sub>3</sub> ).....			0.3	0			
29	Silica (SiO <sub>2</sub> ) Gravimetric.....							
30	Colorimetric.....			9.3	18			
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....			55.0	178			
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....			18.7	19.7			
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....			73.7	198			
34	Sum of Constituents.....			84.0	225			
35	Saturation index.....			-0.15	+0.7			
	Remarks:							

TABLE III—Continued  
**Chemical Analyses of Civic Water Supplies**  
**FRASER RIVER DRAINAGE BASIN**  
*(In parts per million)*

STEVESTON	SUNBURY	SURREY DISTRICT MUNICIPALITY	VANCOUVER			VEDDER CROSSING	No.				
Supplied from Greater Vancouver Water District through Richmond District Municipality	Supplied by Delta District Municipality.	Supplied from Greater Vancouver Water District; White Rock Waterworks Co., Ltd., Crescent Beach Waterworks Ltd., City of Blaine, U.S.A. and Cloverdale, B.C.	Seymour Lake	Capilano River	Coquitlam (River) Lake	A portion supplied with Chilliwack River					
			Raw and finished water			Raw and finished water					
			Spillway at reservoir	Direct from intake in mountains	At lake						
See Vancouver, B.C.	See Delta District Municipality.	See Vancouver, White Rock, Crescent Beach and Cloverdale, B.C.	3391	3423	3438	See Table II. Station No. 22.	1				
			473	474	475		2				
			Aug. 10/49	Aug. 10/49	Aug. 10/49		3				
			27	35	49		4				
			14.0	12.5	17.0		5				
			21.0	24.0 (19.0)	21.0		6				
			..... (3.0)				7				
			7.1 (7.1)	7.5 (7.2)	7.3 (6.9)		8				
			5 (8)	5 (10)	10 (8)		9				
			0.7	0.6	0.9		10				
			.....				11				
			16.4	19.0	12.8		12				
			6.4	7.0	6.4		13				
			23.9	21.8	13.5		14				
			2.2	4.0	1.8		15				
			0.2	0.3	0.2		16				
			.....				17				
			0.15	0.09	0.05		18				
			1.0	0.7	0.9		19				
			0.4	0.3	0.2		20				
			0 (0)	0 (0)	0 (0)		21				
			9.3 (7.3)	8.8 (7.3)	7.6 (7.3)		22				
			2.8	6.5	2.0		23				
			0	0	1.4		24				
			0.05	0.20	0		25				
			Trace	Trace	0		26				
			.....				27				
			4.4	4.0	3.5		28				
			6.3	7.2 (6.0)	5.3		29				
			0	4.0	0		30				
			6.3	11.2	5.3		31				
			15.8	20.4	13.9		32				
			-3.1	-2.4	-3.0		33				
					Vancouver supply is that of Greater Vancouver Water District.						

TABLE III—Concluded  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

No.	Municipality.....	WELLS		WEST VANCOUVER DISTRICT MUNICIPALITY	WHALLEY
		Mosquito Creek	Red Gulch Creek	Supplied from Greater Vancouver Water District	Supplied by Surrey District Municipality
	Source(s).....	Raw and unfinished water			
	Sampling Point.....	From intakes			
1	Laboratory number.....	5023	5024	See Vancouver, B.C.	See Surrey District Municipality
2	Field number.....	808	809		
3	Date of collection.....	Apr. 15/51	Apr. 15/51		
4	Storage period (days).....	5	5		
5	Sampling temperature, °C.....				
6	Test temperature, °C.....	23.5	23.5		
7	Dissolved oxygen.....				
8	Carbon dioxide (CO <sub>2</sub> ).....				
9	pH.....	7.4	7.4		
10	Colour.....	10	5		
11	Turbidity.....	0.5	0.5		
12	Suspended matter, dried at 105°C.....				
13	Suspended matter, ignited at 550°C.....				
14	Residue on evaporation, dried at 105°C.....	51.4	69.8		
15	Ignition loss at 550°C.....	11.0	5.8		
16	Specific conductance (micromhos at 25°C.).....	74.6	120		
17	Calcium (Ca).....	8.9	17.8		
18	Magnesium (Mg.).....	2.9	3.4		
19	Iron (Fe) Total.....				
20	Dissolved.....	0.03	0.01		
21	Sodium (Na).....	1.3	1.5		
22	Potassium (K).....	0	0		
23	Carbonate (CO <sub>3</sub> ).....	0	0		
24	Bicarbonate (HCO <sub>3</sub> ).....	31.7	58.6		
25	Sulphate (SO <sub>4</sub> ).....	9.2	11.6		
26	Chloride (Cl).....	0	0		
27	Fluoride (F).....	0	0		
28	Nitrate (NO <sub>3</sub> ).....	0.4	0		
29	Silica (SiO <sub>2</sub> ) Gravimetric.....				
30	Colorimetric.....	5.9	10		
31	Carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	26.0	48.0		
32	Non-carbonate hardness as CaCO <sub>3</sub> , p.p.m.....	8.2	10.5		
33	Total hardness as CaCO <sub>3</sub> , p.p.m.....	34.2	58.5		
34	Sum of Constituents.....	44.2	73.4		
35	Saturation index.....	-1.6	-1.1		
	Remarks:				



TABLE III—Concluded  
Chemical Analyses of Civic Water Supplies  
FRASER RIVER DRAINAGE BASIN  
(In parts per million)

WHITE ROCK					WILLIAMS LAKE		No.
Gracine Spring	Civic Mixture	Well No. 1	Well No. 2	Well No. 5	Williams Lake		
Raw and finished water					Raw and finished water		
At spring	Town tap	At well	At well	At well	Direct at intake		
4089	4090	4091	4092	4093		4568	1
560	561	558	559	562		733	2
Mar. 20/50	Mar. 20/50	Mar. 20/50	Mar. 20/50	Mar. 20/50	1949	Aug. 21/50	3
12	12	12	12	12		51	4
8.9	11.1	11.7	11.7			20.6	5
21.0	21.0	21.0	21.0	21.0		21.2 (22.8)	6
							7
						(0)	8
						8.5 (8.8)	9
7.7	8.2	8.2	8.2	8.0	8.0	10 (30)	10
0	0	5	5	5	15	0.6	11
0.3	0.2	0.3	0.3	0.5	Slight		12
							13
							14
96.2	159	537	408	153	424	346	15
19.0	12.2	13.0	16.0	12.2	100	91.8	16
129	242	895	681	242		525	17
10.0	23.2	14.6	16.6	24.4	29.8	31.3	18
3.2	8.8	9.0	9.4	9.0	40.2	37.5	19
					0.07		20
0.04	0.03	0.05	0.03	0.02		0.08	21
11.2	16.8	175	123	15.8	as Na { 36.8	32.5	22
1.1	2.7	9.2	7.0	2.5		4.5	23
0	2.4	5.3	3.6	2.4	0	27.4 (30.0)	24
26.4	113	196	168	111	185	314 (300)	25
12.7	14.0	34.4	26.7	11.7	12.6	15.1	26
15.0	16.0	174	124	16.0	0.8	0 (0)	27
0	0.05	0.25	0.10	0.10	0	0.25	28
9.8	0.35	0	Trace	0.4	0.1	0.7	29
12	23	24	23	21	14		30
13	21	25	24	20		11	31
21.6	94.1	73.4	80.1	95.0	239.3	232	32
16.6	0	0	0	2.9	0	0	33
38.2	94.1	73.4	80.1	97.9	239.3	232	34
89.0	161	545	418	157	226	315	35
-1.4	+0.07	+0.06	+0.05	-0.12	+0.5	+1.0	
					Analyses by Dept. Nat. Health and Welfare, Victoria, B.C.		
					Nitrite—0 p.p.m. Fluoride—0.1 p.p.m. (1950)		

**TABLE IV**  
**Municipal Water Supplies Within the Fraser River Drainage Basin**  
Summary of data on area, total population and population served

Region	Approximate area <sup>a</sup>		Estimated total population in thousands	Estimated population in thousands served by organized water systems		Per cent population served in basin (1951)	Per cent population served (1951) with			
	Square miles	Per cent of total province					Soft water	Medium hard water	Hard water	Very hard water
			1951 <sup>b</sup>	1949-50 <sup>c</sup>	1951 <sup>d</sup>					
Lower Fraser River Basin.....	8,520	2.33	644.76	644.1	564.2	87.5	95.6	1.4	3.0	.....
Central Fraser River Basin.....	35,530	9.70	43.66	22.3	23.0	52.6	66.1	13.5	10.4	10.0
Upper Fraser River Basin.....	47,390	12.94	32.91	8.4	8.6	26.1	26.7	73.3	.....	.....
Total basin in Canada.....	91,440	25.0	721.3	674.8	595.8	82.6	93.4	3.0	3.2	0.4
Total province.....	366,255 (359,280 land area)	100.0	1,165.2							

<sup>a</sup> Includes fresh water.

<sup>b</sup> Ninth census of Canada.

<sup>c</sup> Estimated from figures supplied by officials, and from other sources.

<sup>d</sup> Estimated from ninth census of Canada.

**TABLE V**  
**Municipal Water Supplies within the Fraser River Drainage Basin**  
Summary of data on systems, including source, treatment and hardness of waters

Region	Number of municipalities and communities served by organized systems	Number different sources <sup>a</sup>	Source								Per cent of systems using				Per cent of systems using surface waters	Treatment methods, 1950-51		
			Surface waters				Ground waters									None	Chlorination	
			Soft	Med. hard	Hard	Very hard	Soft	Med. hard	Hard	Very hard	Soft	Med. hard	Hard	Very hard			Alone	Additional treatment <sup>c</sup>
Lower Fraser River Basin..	47	13(4)	8(2)	.....	1	.....	.....	4(2)	.....	.....	61.5	30.8	7.7	.....	69.2	11(3)	2	.....
Central Fraser River Basin..	17	14(4)	7(4)	3	1	3	.....	.....	.....	.....	50.0	21.4	7.2	21.4	100	8(3)	4	2
Upper Fraser River Basin..	5	5(1)	3(1)	2	.....	.....	.....	.....	.....	.....	60.0	40.0	.....	.....	100	3(1) <sup>b</sup>	1	1
Total Basin.....	69	32(9)	18(7)	5	2	3	.....	4(2)	.....	.....	56.3	28.1	6.3	9.3	87.5	22(7)	7	3

<sup>a</sup> Figures in brackets refer to number of systems not studied in detail.

<sup>b</sup> One source is naturally filtered.

<sup>c</sup> Additional treatment is natural filtration.

## DISCUSSION

The basin has been divided into three general regions primarily because of the preponderance of population in the lower Fraser River basin or delta area. These regional areas are only rough estimates but do serve to indicate the density of population in each. Continual expansion of the Greater Vancouver Water District, the fact that some communities are only partially served from this District supply, and the shifting and growing population and industrial activity in the area, all tend to outdate statistics in this region.

This is apparent in the rather widely different figures given for population in various communities in 1949 and in the 1951 census. An attempt has been made to arrive at a reasonably accurate estimate using data from several sources, although generally the 1951 census figure, corrected for any known population served outside the incorporated area, has been used in compilation of the tables.

Since most of the data is based on information obtained in 1950 or 1951 no attempt has been made to estimate, by assuming a steady increase from 1941 to 1951, populations for 1949 or 1950; rather, the best data of 1951 have been used. In the preparation of the data of Tables IV and V it has once more been assumed that those systems which were not studied in detail used soft surface water without any treatment. This is a reasonable assumption since these systems are usually very small and creek waters are readily obtained by gravity.

From these tables it is noted that the Fraser River basin which is about 25 per cent of the total area of the province contains about 62 per cent of the provincial population but that about 89 per cent of the basin population or 56 per cent of the provincial population resides in the lower Fraser River basin, mostly concentrated on the delta of the river. Because of the Greater Vancouver Water District, 87.5 per cent of the population in this lower basin region is served with water by organized system, about 96 per cent of those served using soft water. Even though the other portions of the basin have much lower percentage served and generally use harder waters, about 83 per cent of the entire river basin is served with water; 93 per cent of those served using soft water.

The decrease in hardness of surface waters along the Fraser River is indicated somewhat by the data of this Table in that in the upper portion of the basin 73 per cent of those served are served with medium hard water.

The effect of the Greater Vancouver Water District is again noted in Table V in that, while about 47 communities are served in this lower river area, only about 13 systems have different water sources and some of these also use water from the Water District. In this area are the only systems using ground waters, which are medium hard in character. It is estimated that the Greater Vancouver Water District serves about 82 per cent of the population in this region with a soft water.

As in other basins in this province treatment of the water is practically nil, 69 per cent of the systems having no treatment, the remainder being only chlorinated, except for three which have, in addition, natural filtration through gravel beds.

Since the Fraser River itself is not used by any municipality and all other rivers and streams are usually clear, no clarifying treatment other than occasionally coarse screening is required.

## SUMMARY

Most of this basin which contains such a large percentage of the province's population and industry has available an adequate supply of soft water cheaply distributed with little or no treatment. Even in those areas where the main rivers are such as to require at least some clarifying treatment there is usually readily available by gravity clear, soft to medium hard, mountain streams requiring no treatment.

Pollution of the basin's many water sources is inappreciable as evidenced by the lack of chlorination. Since much of the basin is only sparsely settled, industrial activity is small and water sources are in unsettled mountainous regions, it is not likely that pollution will be at all serious for many years. In the more heavily-populated areas steps have already been taken to protect from pollution the headwaters of the water sources, which are usually in nearby mountainous areas.

Water supply in this basin is adequate for a much larger population and extensive industrial use. For most industrial uses the only treatment required would be clarification of turbid waters or prevention of corrosion by the soft, oxygen-saturated waters, typical of this province and other mountainous regions.





# APPENDIX A

## SAMPLING LOCATIONS OF SURFACE WATERS

Station No.	PAGE
39. Adams River near Squilax.....	40
17. Alouette River near Haney.....	26
52. Bonaparte River at Cache Creek.....	44
55. Bridge River near Lillooet.....	44
70. Burns Lake near Burns Lake.....	54
20. Cannell Lake near Mission City.....	28
64. Chilako River near Prince George.....	50
58. Chilcotin River near Alexis Creek.....	46
59. Chilko River near Redstone.....	46
22. Chilliwack River at Vedder Crossing.....	28
36. Clearwater River near Clearwater.....	38
51. Coldwater River at Merritt.....	44
26. Coquihalla River at Hope.....	30
62. Cottonwood River near Quesnel.....	48
23. Cultus Lake at Cultus Lake.....	30
53. Deadman River near Savona.....	44
46. Eagle River near Sicamous.....	42
47. Eagle River near Malakiva.....	42
45. East Canoe Creek near Salmon Arm.....	30
24. Elk River near Chilliwack.....	
Fraser River.....	14
1. (1) at Haney.....	14
2. (2) at Mission City.....	16
3. (3) at Rosedale.....	18
4. (4) at Hope.....	20
5. (5) at Spuzzum.....	20
6. (6) at Boston Bar.....	20
7. (7) at Lillooet.....	20
8. (8) near Williams Lake.....	22
9. (9) at Macalister.....	22
10. (10) at Quesnel.....	24
11. (11) below Prince George.....	24
12. (12) above Prince George.....	26
13. (13) near Tête Jaune Cache.....	
14. (14) near Mount Robson.....	30
25. Harrison River near Harrison Mills.....	48
60. Horsefly River at Horsefly.....	32
28. Lytton Creek at Lytton.....	42
42. Mara Lake near Mara.....	42
43. Mara Lake near Sicamous.....	
Nechako River.....	50
63. (1) at Prince George.....	50
65. (2) at Vanderhoof.....	52
67. (3) at Fort Fraser.....	42
49. Nicola River at mouth.....	44
50. Nicola River near Nicola.....	52
68. Northerly River near Fort Fraser.....	42
48. North Fork Eagle River (Perry River) near Craigellachie.....	

# APPENDIX A—Concluded

## SAMPLING LOCATIONS OF SURFACE WATERS—Concluded

<i>Station No.</i>		<i>PAGE</i>
	North Thompson River	
33.	(1) at Rayleigh.....	36
34.	(2) at Barriere.....	38
35.	(3) at Clearwater.....	38
16.	Pitt River near mouth.....	26
61.	Quesnel River near Quesnel.....	48
37.	Raft River near Clearwater.....	38
44.	Salmon River near Salmon Arm.....	42
56.	San José River (Lac la Hache) near Wright.....	44
27.	Schkam Creek at Hope.....	32
54.	Seton Lake near Lillooet.....	44
40.	Shuswap Lake at Salmon Arm.....	40
41.	Shuswap River near Enderby.....	40
38.	South Thompson River at Chase.....	40
18.	Stave River at Stave Falls.....	26
19.	Stave River near Ruskin.....	26
69.	Stellako River near Fort Fraser.....	52
66.	Stuart River at Fort St. James.....	50
21.	Sumas River near Kilgard.....	28
	Thompson River	
29.	(1) at Spences Bridge.....	32
30.	(2) at Ashcroft.....	32
31.	(3) near Savona.....	34
32.	(4) below Kamloops.....	34
57.	Williams Lake at Williams Lake.....	44
71.	Willow River at Willow River.....	54
15.	Yellowhead Lake near Lucerne.....	26

# APPENDIX B CIVIC WATER SUPPLIES IN THE FRASER RIVER DRAINAGE BASIN

	DATA PAGE	ANALYSIS PAGE		DATA PAGE	ANALYSIS PAGE
Abbotsford <sup>a</sup> .....	58	74	Maillardville.....	65	78
Armstrong <sup>a</sup> .....	58	74	Maple Ridge District Municipality	65	79
Ashcroft <sup>a</sup> .....	59	74	Matsqui District Municipality <sup>a</sup> ....	64	79
Beach Grove <sup>a</sup> .....	59	74	McBride <sup>a</sup> .....	64	79
Bralorne <sup>a</sup> .....	59	74	Merritt <sup>a</sup> .....	65	79
Bridgeport.....	58	74	Mission City <sup>a</sup> .....	65	79
Brighouse.....	58	74	Mission District Municipality.....	65	79
Burkeville.....	59	74	Newton Station.....	66	80
Burns Lake <sup>a</sup> .....	59	75	New Westminster.....	66	80
Burnaby.....	59	75	North Kamloops.....	67	80
Burquitlam.....	60	75	North Vancouver <sup>a</sup> .....	67	80
Canoe.....	60	75	North Vancouver Dis. Municipality	67	80
Chase <sup>a</sup> .....	61	—	Pitt Meadows District Municipality	66	81
Chilliwack <sup>a</sup> .....	61	75	Port Coquitlam.....	67	81
Chilliwack District Municipality..	61	75	Port Mann.....	66	81
Clinton <sup>a</sup> .....	60	75	Port Moody.....	67	81
Cloverdale <sup>a</sup> .....	60	76	Prince George <sup>a</sup> .....	67	81
Coquitlam District Municipality...	61	76	Quesnel <sup>a</sup> .....	68	81
Crescent Beach <sup>a</sup> .....	61	76	Richmond District Municipality...	69	82
Delta District Municipality.....	61	76	Rosedale.....	68	82
East Richmond.....	62	76	Salmon Arm <sup>a</sup> .....	69	82
Eburne.....	62	76	Salmon Arm District Municipality.	69	82
Enderby <sup>a</sup> .....	62	77	Sardis.....	68	82
Essondale.....	63	77	South Westminster.....	68	82
Fraser Mills District Municipality..	63	77	Steveston.....	69	83
Gold Bridge <sup>a</sup> .....	63	77	Sunbury.....	69	83
Haney.....	62	77	Surrey District Municipality.....	69	83
Harrison Hot Springs <sup>a</sup> .....	62	77	Vancouver <sup>a</sup> .....	70	83
Hope <sup>a</sup> .....	63	77	Vedder Crossing <sup>a</sup> .....	70	83
Ioco <sup>a</sup> .....	63	—	Walhachin <sup>a</sup> .....	71	—
Kamloops <sup>a</sup> .....	63	78	Wells <sup>a</sup> .....	71	84
Kennedy.....	64	78	West Vancouver Dis. Municipality	71	84
Ladner.....	64	78	Whalley.....	72	84
Lillooet <sup>a</sup> .....	64	78	White Rock <sup>a</sup> .....	72	85
Lytton <sup>a</sup> .....	65	78	Williams Lake <sup>a</sup> .....	72	85

<sup>a</sup>Communities known to have separate or different sources of supply.  
<sup>a</sup>No organized water system.





THE LIMITS OF THE ARBITRARY SUBDIVISIONS OF THE FRASER RIVER DRAINAGE BASIN AS SHOWN ON THIS MAP ARE INACCURATE. THE NORTHERN LIMIT OF THE CENTRAL BASIN LIES ALONG LATITUDE 52° 35'. THE SOUTHERN LIMIT FOLLOWS THE BOUNDARY AND NORTH OF THE GLOUCESTER RIVER, CROSSING THE FRASER RIVER SOUTH OF LYTON AND THENCE TO THE HEADWATERS OF THE COQUILZINE RIVER.

REF. PP. 96 AND 97 OF C.A.S.I.

CANADA  
DEPARTMENT OF  
MINES AND TECHNICAL SURVEYS

MINES BRANCH

FIGURE 2  
Civic water supplies  
and locations of sampling stations,  
Fraser River drainage basin,  
British Columbia

Scale: 1 Inch to 30 Miles 1:360,000



LEGEND


Civic Supplies	Sampling Stations
Total Hardness as ppm CaCO <sub>3</sub>	Daily Sampling
Unknown	Monthly Sampling
0-50	Spot Sampling
51-120	
121-180	
Over 180	
Boundary of drainage basin	
Municipality boundary (on inset map)	



## NOTE

THE LIMITS OF THE ARBITRARY SUBDIVISIONS OF THE FRASER RIVER DRAINAGE BASIN AS SHOWN ON THIS MAP ARE INCORRECT.

THE NORTHERN LIMIT OF THE CENTRAL BASIN LIES ALONG LATITUDE 52° 30'; THE SOUTHERN LIMIT FOLLOWS THE HEIGHT OF LAND NORTH OF THE LILLOOET RIVER, CROSSING THE FRASER RIVER SOUTH OF LYTTON AND THENCE TO THE HEADWATERS OF THE COQUIHALLA RIVER.





71  
DEPARTMENT OF GEOLOGICAL SCIENCES C.L.  
Author Thomas, J.F.J. (Dept. Mines & Tech. Surveys)  
Water Survey Report No. 6. Fraser River  
Title Drainage Basin, 1950-51.

622  
C16  
17-I  
6



